arXiv:1105.1544v1 [math.DG] 8 May 2011

EXTREMAL OF LOG SOBOLEV INEQUALITY AND W ENTROPY ON NONCOMPACT MANIFOLDS

QI S. ZHANG

Abstract. Let $\mathbf{M}$ be a complete, connected noncompact manifold with bounded geometry. Under a condition near infinity, we prove that the Log Sobolev functional (1.1) has an extremal function decaying exponentially near infinity. We also prove that an extremal function may not exist if the condition is violated. This result has the following consequences. 1. It seems to give the first example of connected, complete manifolds with bounded geometry where a standard Log Sobolev inequality does not have an extremal. 2. It gives a negative answer to the open question on the existence of extremal of Perelman's W entropy in the noncompact case, which was stipulated by Perelman [P] p9, 3.2 Remark. 3. It helps to prove, in some cases, that noncompact shrinking breathers of Ricci flow are gradient shrinking solitons.

Contents

1. Introduction 1
2. Proof of Theorem 1.1 (a), the existence part 4
3. Proof of the theorem 1.1 (b), the nonexistence part 15
4. W entropy and a no breather result for noncompact Ricci flow 34
References 43

1. Introduction

The main purpose of the paper is to give a counter example to the old question on existence of extremals of a standard Log Sobolev inequality (or its recent reincarnation in the form of Perelman's W entropy) on noncompact manifolds with bounded geometry. We also prove existence of extremal under an extra condition. Finding extremal of useful functionals is an useful problem in mathematical analysis. For instance there is a vast literature devoted to the study of ground state eigenvalues and eigenfunctions which are extremal of the Dirichlet functional. The Log Sobolev functional (1.1) seems to be a mild nonlinear perturbation of the Dirichlet functional. Indeed, they share a common property i.e. there exist extremal functions for both functionals on compact domains or compact manifolds. However in the noncompact case the similarity stops. For instance in $\mathbf{R^n}$, it is well known that the Dirichlet functional does not have an extremal or L^2 eigenfunction. In contrast the Gaussian functions are extremals of the Log Sobolev functional. Over the years, Log Sobolev inequality has found many applications in various branches of mathematics and physics. See for example the papers Gross [G], [G2], Federbush [F],

---

Date: May 2011.

Bakry and Émery [BE], Bakry and Ledoux [BL], Diaconis and Saloff-Coste [DS] and Otto and Villani [OV]. A more recent application was discovered by Perelman [P] where he introduced the fundamental W entropy (4.1) and used it as a key analytic tool to prove the Poincaré conjecture. The W entropy is just the Log Sobolev functional (1.1) scaled with certain time dependent parameter. For the Log Sobolev functional, the existence problem of extremal functions in the compact case was solved by O. Rothaus [Rot] 30 years ago. However, in the noncompact case, the problem is wide open. There has been no counter example or general existence result for connected, noncompact manifolds with bounded geometry. We should mention that if one drops the connectedness, then it is easy to construct a manifold with infinitely many disconnected components, such that the Log Sobolev functional does not have an extremal. See the example at the beginning of Section 3 e.g. Also if the manifold is homogeneous such as $\mathbf{R^n}$, one can use symmetrization or translation or group action to prove existence of an extremal.

In addition to being an interesting problem in its own right, the study of Log Sobolev inequality or W entropy in the noncompact setting is also important to Ricci flow. One reason is that many of the more interesting singularity models are noncompact, even when the Ricci flow under consideration is compact. One such example in the three dimensional case is the round neck $S^2 \times R$, which is a typical singularity model. Using the existence of extremals of his W entropy, Perelman [P] proved a no breather theorem stating that shrinking breathers of Ricci flows on compact manifolds are shrinking gradient solitons. Recently, in the case $(\mathbf{M}, g)$ is a noncompact gradient shrinking soltion, Carrillo and Ni [CN] proved that potential functions are extremals for W the entropy.

On p9, 3.2 Remark of the same paper, Perelman also wrote

"Of course, this argument requires the existence of minimizer, and justifications of the integration by parts; this is easy if M is closed, but can also be done with more efforts on some complete M, ..."

However, it is not known so far if the W entropy always has an extremal for all noncompact manifolds which are reasonably nice, such as those connected ones with bounded geometry. The main theorem of the paper (Theorem 1.1 or Theorem 4.1) shows that on noncompact manifolds, the Log Sobolev functional or the W entropy has an extremal function under a condition near infinity; it also shows that an extremal function may not exist if the condition is violated, giving a negative answer to the above question stipulated by Perelman. As another application we partially extend Perelman's no breather theorem to the noncompact case. See Section 4 below.

In order to state the result precisely, we first introduce a number of basic assumptions and notations.

Basic assumptions. In this paper, unless stated otherwise, we assume the n dimensional Riemannian manifold $\mathbf{M}$ with metric g is a complete noncompact manifold with bounded geometry which means:

1. there exists a positive constant α such that

$$|Rm| \le \alpha$$

where Rm is the curvature tensor and $|Rm|$ is the maximum norm of Rm under g.

2. there exists a positive constant β such that, for all $x \in \mathbf{M}$,

$$|B(x, 1)|_g \ge \beta.$$

Here $B(x,1)$ is the geodesic ball of radius 1, centered at x; and $|B(x,1)|_g$ is the volume of $B(x,1)$ under the metric g.

It is well known that assumptions 1 and 2 imply that the injectivity radius of $\mathbf{M}$ is bounded from below by a positive constant. See [CGT] and [CLY] e.g.

We will use the following notations throughout the paper. g_{ij}, R_{ij} will be the metric and Ricci curvature; R is the scalar curvature; ∇, Δ the corresponding gradient and Laplace-Beltrami operator; dg is the volume element; c, C with or without index denote generic positive constant that may change from line to line.

The main Log Sobolev inequality that we deal with in this paper is just the usual one perturbed by the scalar curvature of the manifold. i.e. there exist positive constant a and another constant $c = c(a, \mathbf{M}, g)$ such that, for $v \in C_0^\infty(\mathbf{M})$ and $\|v\|_{L^2(\mathbf{M})} = 1$,

$$\int_{\mathbf{M}} v^2 \ln v^2 dg \le a \int_{\mathbf{M}} (4|\nabla v|^2 + Rv^2) dg + c(a, \mathbf{M}, g).$$

The functional associated with the Log Sobolev inequality when $a = 1$ is

$$L(v,g) \equiv \int_{\mathbf{M}} (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg, \qquad v \in W^{1,2}(\mathbf{M}). \tag{1.1}$$

One reason for involving the scalar curvature is, after scaling the functional by certain time dependent factor and coupled with Ricci flow, it becomes Perelman's W entropy [P], which is a fundamental quantity for Ricci flow. This relation is shown in (4.2). The existence and nonexistence of extremal of the Log Sobolev functional depends on two quantities given in the definition below. The first one is just the best Log Sobolev constant or the infimum of the functional in (1.1). The second one is the best Log Sobolev constant at infinity. The concept is motivated by P.L. Lions' concentration compactness principle [L].

Definition 1.1. *Let* $(\mathbf{M}, g)$ *be a complete noncompact manifold with bounded geometry.*

The best Log Sobolev constant of $(\mathbf{M}, g)$ *is the quantity*

$$\lambda = \lambda(\mathbf{M}) = \lambda(\mathbf{M}, g) = \inf\{\int_{\mathbf{M}} (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg \quad | \, v \in C_0^\infty(\mathbf{M}), \quad \|v\|_{L^2(\mathbf{M}) = 1}\}.$$

The best Log Sobolev constant of $(\mathbf{M}, g)$ *at infinity is the quantity*

$$\lambda_\infty = \lambda_\infty(\mathbf{M}, g) = \lim_{r \to \infty} \inf\{\int_{\mathbf{M}-B(0,r)} (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg \quad | \quad v \in C_0^\infty(\mathbf{M} - B(0,r)), \quad \|v\|_{L^2(\mathbf{M}-B(0,r)) = 1}\}.$$

Let D *be a domain in* $\mathbf{M}$*. The best Log Sobolev constant of* D *is the quantity*

$$\lambda = \lambda(D) = \lambda(D, g) = \inf\{\int_D (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg \quad | \, v \in C_0^\infty(D), \quad \|v\|_{L^2(D) = 1}\}.$$

Definition 1.2. *(extremal) Suppose* $\lambda = \lambda(\mathbf{M}, g)$ *is a finite number. A function* $v \in W^{1,2}(\mathbf{M})$ *is called an extremal of the Log Sobolev functional (1.1) if* $\|v\|_{L^2(\mathbf{M})} = 1$ *and*

$$\int_{\mathbf{M}} (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg = \lambda$$

The main result of the paper is the following theorem, or equivalently Theorem 4.1 in Section 4.

Theorem 1.1. *(a). Let* $\mathbf{M}$ *be a complete, connected noncompact manifold with bounded geometry. Suppose* $\lambda < \lambda_\infty$*, then there exists a smooth extremal* v *for the Log Sobolev functional in (1.1). Also, there exist positive constants* $a, A > 0$ *and a point* $0 \in \mathbf{M}$ *such that*

$$v(x) \le A e^{-ad^2(x,0)}.$$

(b). There exists a complete, connected noncompact manifold with bounded geometry such that $\lambda = \lambda_\infty$*, but the Log Sobolev functional in (1.1) does not have an extremal.*

Remark. Manifolds satisfying the condition $\lambda < \lambda_\infty$ are quite common. For example, suppose $\mathbf{M}$ is asymptotically Euclidean, then $\lambda_\infty = \lambda(\mathbf{R}^n) = \frac{n}{2}\ln(4\pi) - n$. If there exists a compact domain $D \subset \mathbf{M}$ such that $\lambda(D) < \frac{n}{2}\ln(4\pi) - n$. Then

$$\lambda \le \lambda(D) < \lambda_\infty.$$

It is easy to construct a domain such that $\lambda(D)$ is arbitrarily negative. One example is the scaled flat torus $h^2(S^1 \times S^1) \times S^1$ when the scaling factor $h \to 0$. See Lemma 3.6.

Even though the Log Sobolev functional in the theorem contains the scalar curvature R, the result still holds if one deletes the scalar curvature. The proof only requires minor adjustment.

The rest of the paper is organized as follows. Theorem 1.1 (a) and (b) will be proven in Sections 2 and 3 respectively. Applications on the W entropy will be given in Section 4.

2. Proof of Theorem 1.1 (a), the existence part

The proof of the theorem relies on the study of the Euler-Langrange equation of the Log Sobolev functional:

$$4\Delta v - Rv + 2v \ln v + \lambda v = 0. \tag{2.1}$$

When λ is the best Log Sobolev constant, this is the equation satisfied by the extremal. Sometimes we also need to deal with subsolutions to this equation. A function $v \in W^{1,2}_{loc}(\mathbf{M})$ is called a subsolution to (2.1) if it satisfies the following inequality in the weak sense:

$$4\Delta v - Rv + 2v \ln v + \lambda v \ge 0, \qquad \text{in} \qquad \mathbf{M}. \tag{2.2}$$

i.e., for any nonnegative, compactly supported test function ψ, we have

$$\lambda \int_{\mathbf{M}} v\psi dg \ge \int_{\mathbf{M}} (4\nabla v \nabla \psi + Rv\psi - 2\psi v \ln v) dg.$$

We will need a number of lemmas before proving the theorem. The first lemma is a mean value type inequality for subsolutions of the above equation

Lemma 2.1. *(a). Suppose* v *is a bounded subsolution to the equation (2.1) in the ball* $B(m, 2) \subset \mathbf{M}$ *such that* $\|v\|_{L^2(B(m,2))} \le 1$*. Here* $m \in \mathbf{M}$ *which has bounded geometry. Then there exists a positive constant* $C = C(n, \alpha, \beta, \lambda)$ *such that*

$$\sup_{B(m,1)} v^2 \le C \int_{B(m,2)} v^2 dg.$$

(b). Moreover, if v is a bounded solution to (2.1) in the ball $B(m, 2) \subset \mathbf{M}$ such that $\|v\|_{L^2(B(m,2))} \le 1$, then there exists a positive constant $C = C(n, \alpha, \beta, \lambda)$ such that the gradient bound holds:

$$\sup_{B(m,1/2)} |\nabla v|^2 \le C \int_{B(m,1)} v^2 dg.$$

Proof.

Part (a). This part of the lemma and its proof is similar to that of Lemma 8.2.1 in [Z] where the underlying manifold is an ϵ horn. The proof relies on Moser's iteration and standard Sobolev inequality and takes advantage of the slow growth of $\ln v$ when v is large.

Given any $p \ge 1$, it is easy to see that

$$-4\Delta v^p + pRv^p \le 2pv^p \ln v + p|\lambda| v^p. \tag{2.3}$$

We select a smooth cut off function ϕ supported in $B(m, 2)$. Writing $w = v^p$ and using $w\phi^2$ as a test function in (2.3), we deduce

$$4 \int \nabla(w\phi^2)\nabla w dg + p \int R(w\phi)^2 dg \le 2p \int (w\phi)^2 \ln v dg + p \int |\lambda|(w\phi)^2 dg.$$

By the bound on the curvature tensor $|Rm| \le \alpha$, we deduce

$$4 \int \nabla(w\phi^2)\nabla w dg \le p \int (w\phi)^2 \ln v^2 dg + (C\alpha + |\lambda|)p \int (w\phi)^2 dg,$$

which induces, after integration by parts,

$$4 \int |\nabla(w\phi)|^2 dg \le 4 \int |\nabla\phi|^2 w^2 dg + (C\alpha p + |\lambda|) \int (w\phi)^2 dg + p \int (w\phi)^2 \ln v^2 dg. \tag{2.4}$$

We need to dominate the last term in (2.4) by the left hand side of (2.4). For one positive number a to be chosen later, it is clear that

$$\ln v^2 \le v^{2a} + c(a).$$

Hence for any fixed $q > n/2$, the Hölder inequality implies

$$\begin{aligned} p \int (w\phi)^2 \ln v^2 dg &\le p \int (w\phi)^2 v^{2a} dg + pc(a) \int (w\phi)^2 dg \\ &\le p \left(\int v^{2aq} dg \right)^{1/q} \left(\int (w\phi)^{2q/(q-1)} dg \right)^{(q-1)/q} + pc(a) \int (w\phi)^2 dg. \end{aligned}$$

We take $a = 1/q$ so that $2aq = 2$. Since the L^2 norm of u is less than 1 by assumption, the above implies

$$p \int (w\phi)^2 \ln v^2 dg \le p \left(\int (w\phi)^{2q/(q-1)} dg \right)^{(q-1)/q} + pc(a) \int (w\phi)^2 dg.$$

By interpolation inequality (see p84 [HL] e.g.), it holds, for any $b > 0$,

$$\left(\int (w\phi)^{2q/(q-1)} dg \right)^{(q-1)/q} \le b \left(\int (w\phi)^{2n/(n-2)} dg \right)^{(n-2)/n} + c(n, q) b^{-n/(2q-n)} \int (w\phi)^2 dg.$$

Therefore

(2.5)
$$p\int (w\phi)^2 \ln v^2 dg \le pb\left(\int (w\phi)^{2n/(n-2)} dg\right)^{(n-2)/n} + Cpb^{-n/(2q-n)} \int (w\phi)^2 dg + pC\int (w\phi)^2 dg.$$

Since the manifold $\mathbf{M}$ has bounded geometry, it is well known ([Au],[Heb], [HV] and [Sa] e.g.) that a standard Sobolev inequality holds, i.e. there exist positive constants S_0 depending on α, β, n such that

$$S_0\left(\int (w\phi)^{2n/(n-2)} dg\right)^{(n-2)/n} \le \int |\nabla (w\phi)|^2 dg + \int (w\phi)^2 dg.$$

This and (2.4) imply

(2.6)
$$S_0\left(\int (w\phi)^{2n/(n-2)} dg\right)^{(n-2)/n} \le \int |\nabla \phi|^2 w^2 dg + (C\alpha p + |\lambda| + 1)\int (w\phi)^2 dg + p\int (w\phi)^2 \ln v^2 dg.$$

Substituting (2.5) to the right hand side of (2.6), we arrive at

$$\begin{aligned} S_0\left(\int (w\phi)^{2n/(n-2)} dg\right)^{(n-2)/n} &\le 4\int |\nabla \phi|^2 w^2 dg + pb\left(\int (w\phi)^{2n/(n-2)} dg\right)^{(n-2)/n} \\ &\quad + c(n,q) pb^{-n/(2q-n)} \int (w\phi)^2 dg + pc(a)\int (w\phi)^2 dg. \end{aligned}$$

Take b so that $pb = S_0/2$. It is clear that exist positive constant $c = c(S_0, n, q)$ and $p_0 = p_0(n, q)$ such that

$$\left(\int (w\phi)^{2n/(n-2)} dg\right)^{(n-2)/n} \le c(p+1)^{p_0} \int (|\nabla \phi|^2 + 1) w^2 dg. \tag{2.7}$$

From here one can use standard Moser's iteration to prove the desired bound. We briefly sketch the main steps. Let $\xi_k = \xi_k(s)$, $k = 0, 1, 2, \ldots$, be a smooth one variable function such that $\xi_k(s) = 1$ when $s \in [0, 1 + (1/2^{k+1})]$; $0 \le \xi_k(s) \le 1$, when $s \in [1 + (1/2^{k+1}), 1 + (1/2^k)]$ and $\xi_k(s) = 0$, when $s \in [1 + (1/2^k), \infty)$. We also require that $|\xi'(s)| \le c/2^k$. Set the test function $\phi_k = \xi_k(d(x, m))$. Then it is clear that

$$|\nabla \phi_k| \le \frac{c}{2^k}. \tag{2.8}$$

By (2.7) and (2.8)

$$\left(\int_{B(m, 1+(1/2^{k+1}))} w^{2n/(n-2)} dg\right)^{(n-2)/n} \le \frac{C}{2^{2k}} (p+1)^{p_0} \int_{B(m, 1+(1/2^k))} w^2 dg. \tag{2.9}$$

Recall that $w = v^p$. We iterate (2.9) with $p = (n/(n-2))^k$, $k = 0, 1, 2, \ldots$ Following Moser, we get

$$\sup_{B(m,1)} v^2 \le C\int_{B(m,2)} v^2 dg.$$

This proves part (a) of the lemma.

Part (b). By standard computation, in local orthonormal system, we have

$$\Delta|\nabla v|^2 = 2\Sigma_{i,j} v_{ij}^2 + 2\Sigma_i (\Delta v)_i v_i + 4R_{ij} v_i v_j.$$

Here v_i is the covariant derivative of v and R_{ij} is the Ricci curvature. Since v is a solution to (2.1), we know that

$$(\Delta v)_i v_i = \frac{1}{4}(Rv - 2v \ln v - \lambda v)_i v_i = \frac{1}{4}(R_i v v_i + R v_i^2 - 2v_i^2 \ln v - 2v_i^2 - \lambda v_i^2).$$

Since , by part (a), $v \le C$ in $B(m, 1)$, we have $-\ln v \ge -\ln C$. Hence there exists a positive constant C such that

$$\Delta|\nabla v|^2 \ge -C(|\nabla v|^2 + v^2)$$

in the ball $B(m, 1)$. From here, we can use Moser's iteration for standard Laplacian to conclude that

$$\sup_{B(m,1/2)} |\nabla v|^2 \le C \int_{B(m,2r/3)} (|\nabla v|^2 + v^2) dg \le C \int_{B(m,r)} v^2 dg.$$

□

The next lemma shows that interior maximum value of a positive solution of equation (2.1) in a ball has a positive lower bound independent of the ball. This property in case of compact manifolds was already observed in Section 17.2 of [CCGGIIKLLN3].

Lemma 2.2. *Let v be a smooth positive solution of equation (2.1) in the ball $B(0, r) \subset \mathbf{M}$ such that $v = 0$ on $\partial B(0, r)$. Here 0 is a point in $\mathbf{M}$ and $r > 0$. Then*

$$\sup_{B(0,r)} v \ge e^{(\inf R - \lambda)/2}.$$

i.e. the maximum value of v is bounded from below by a positive constant depending only on λ and the lower bound of the scalar curvature.

Proof.

Since v is 0 at the boundary, clearly the maximum of v is reached at some point x_0 in the interior of the ball $B(0, r)$. Hence $\Delta v(x_0) \le 0$, which implies, by equation (2.1),

$$-R(x_0)v(x_0) + 2v(x_0) \ln v(x_0) + \lambda v(x_0) \ge 0.$$

From this, the lemma follows. □

Lemma 2.3. *Let v be a bounded subsolution to (2.1) on $\mathbf{M}$ such that $\|v\|_{L^2(\mathbf{M})} \le 1$. Let 0 be a reference point on $\mathbf{M}$. Then there exist positive numbers r_0, a and A, which may depend on α, β and the location of the reference point such that*

$$v(x) \le A e^{-a d^2(x,0)}, \qquad when \qquad d(x, 0) \ge r_0.$$

Proof.

Recall from Lemma 2.1 that there exists a constant $C > 0$ such that

$$v^2(x) \le C \int_{B(x,2)} v^2 dg, \qquad x \in \mathbf{M}.$$

This infers

$$-2 \ln v(x) \ge -\ln C - \ln \int_{B(x,2)} v^2 dg.$$

Since $\int_{\mathbf{M}} v^2 dg \le 1$, we know that

$$\lim_{d(x,0)\to\infty} \int_{B(x,1)} v^2 dg = 0.$$

Therefore $-\ln v(x) \to +\infty$ when $d(x,0) \to \infty$. Thus, there exists $r_0 > 0$, such that, when $d(x,0) \ge r_0$, we have

$$R(x) - \ln v(x) - \lambda \ge 0, \qquad \text{and} \quad v(x) \le e^{-1} \tag{2.10}$$

Substituting this to (2.2), we deduce,

$$4\Delta v(x) + v(x) \ln v(x) \ge v(x)(R(x) - \ln v(x) - \lambda) \ge 0.$$

Hence, when $d(x,0) \ge r_0$, we have

$$4\Delta v(x) + v(x) \ln v(x) \ge 0, \qquad \text{and} \quad v(x) \le e^{-1}. \tag{2.11}$$

Next we compare v with a model function

$$J = J(x) = e^{-aL^2(x) + ar_0^2 - 1}. \tag{2.12}$$

Here $a > 0$ is to be decided later; $L = L(x)$ is a smooth function on $\mathbf{M}$, which satisfies

$$\begin{aligned} &|\nabla L(x)| \le C_1, \qquad |\nabla^2 L(x)| \le C_1, \qquad x \in \mathbf{M}, \\ &C_1^{-1} L(x) \le d(x,0) \le C_1 L(x), \qquad d(x,0) \ge r_0. \end{aligned}$$

Under our assumption of bounded geometry, it is well known that such a function exists. For instance, let $\eta \ge 0$ be a smooth function in $C_0^\infty(\mathbf{R}^n)$, supported in a ball centered at the origin, whose radius is less than the injectivity radius of $\mathbf{M}$. If also $\|\eta\|_{L^1(\mathbf{R}^n)} = 1$, then

$$L(x) = \int_{\mathbf{R}^n} \eta(w)[d(0, exp_x(w)) + 1] dw \tag{2.13}$$

satisfies the above requirements. See also the proof of Proposition 19.37 in [CCGGIIKLLN3], e.g. Since $d(x,0)$ and $L(x)$ are comparable when they are large, by (2.11), we can choose r_0 sufficiently large so that

$$4\Delta v(x) + v(x) \ln v(x) \ge 0, \qquad \text{and} \quad v(x) \le e^{-1} \tag{2.14}$$

when $L(x) \ge r_0$.

By direct computation

$$\begin{aligned} \Delta J &= J[4a^2 |\nabla L|^2 L^2 - 2aL\Delta L - 2a|\nabla L|^2], \\ J \ln J &= J(-aL^2 + ar_0^2 - 1). \end{aligned}$$

Hence

$$\begin{aligned} 4\Delta J + J \ln J &= J[16a^2 |\nabla L|^2 L^2 - 8aL\Delta L - 8a|\nabla L|^2 - aL^2 + ar_0^2 - 1] \\ &\le J[16a^2 C_1^2 L^2 + 8aC_1 L - aL^2 + ar_0^2 - 1]. \end{aligned}$$

This implies, for some $C_2 > 0$,

$$4\Delta J + J \ln J \le J[C_2 a^2 L^2 - aL^2 + ar_0^2 - (1/2)].$$

We take $a = \min\{\frac{1}{C_2}, \frac{1}{\sqrt{2C_2 r_0^2}}\}$. Then

$$4\Delta J + J \ln J \le 0$$

when $L(x) \geq r_0$ and $J(x) = e^{-1}$ when $L(x) = r_0$. This and (2.14) show that

$$\begin{cases} 4\Delta(J-v) + J\ln J - v\ln v \leq 0, & \text{if} \quad L(x) \geq r_0, \\ J(x) \leq e^{-1}, \qquad v(x) \leq e^{-1}, & \text{if} \quad L(x) \geq r_0 \\ (J-v)(x) \geq 0, & \text{if} \quad L(x) = r_0, \\ (J-v)(x) \to 0, & \text{if} \quad L(x) \to \infty, \end{cases}$$

Since $J(x), v(x) \leq e^{-1}$, by the mean value theorem, there exists a function $f = f(J(x), v(x))$, $0 < f \leq e^{-1}$ such that

$$J(x)\ln J(x) - v(x)\ln v(x) = (\ln f + 1)(J(x) - v(x)).$$

Observe that

$$\ln f + 1 \leq lne^{-1} + 1 \leq 0, \qquad \text{when} \quad L(x) \geq r_0.$$

Therefore we can apply the standard maximum principle for the elliptic inequality on

$$4\Delta(J-v)(x) + (\ln f + 1)(J-v)(x) \leq 0, \qquad \text{when} \quad L(x) \geq r_0$$

to conclude that

$$v(x) \leq J(x) = e^{-aL^2(x) + ar_0^2 - 1}, \qquad \text{when} \quad L(x) \geq r_0.$$

Since $L(x)$ and $d(x, 0)$ are comparable when they are large, we have proven the lemma by making a smaller if necessary. □

Lemma 2.4. *Let $(\mathbf{M}, g)$ be a complete noncompact manifold with bounded geometry. Let $v \in W^{1,2}(\mathbf{M})$, $\|v\|_{L^2(\mathbf{M})} = 1$ be a bounded sub-solution of (2.1) i.e.*

$$4\Delta v - Rv + 2v\ln v + \lambda v \geq 0.$$

Here λ is a constant. Let D be a bounded domain in $\mathbf{M}$ and define

$$\lambda(D) = \inf\{\int (4|\nabla v|^2 + Rv^2 - v^2\ln v^2)dg \mid v \in C_0^\infty(D), \ \|v\|_2 = 1\}, \tag{2.15}$$

For any smooth cut-off function $\eta \in C_0^\infty(D)$, $0 \leq \eta \leq 1$, it holds

$$\lambda(D)\int (v\eta)^2 dg \leq \lambda \int (v\eta)^2 dg + 4\int v^2|\nabla\eta|^2 dg - \int (v\eta)^2 \ln \eta^2 dg.$$

Proof.

Since $\eta v/\|\eta v\|_2 \in C_0^\infty(D)$ and its L^2 norm is 1, we have, by definition,

$$\lambda(D) \leq \int \left[4\frac{|\nabla(\eta v)|^2}{\|\eta v\|_2^2} + R\frac{(\eta v)^2}{\|\eta v\|_2^2} - \frac{(\eta v)^2}{\|\eta v\|_2^2}\ln\frac{(\eta v)^2}{\|\eta v\|_2^2}\right] dg.$$

This implies

$$\lambda(D)\|\eta v\|_2^2 \leq \int \left[4|\nabla(\eta v)|^2 + R(\eta v)^2 - (\eta v)^2\ln(\eta v)^2\right] dg + \|\eta v\|_2^2 \ln\|\eta v\|_2^2. \tag{2.16}$$

On the other hand, v satisfies

$$4\Delta v - Rv + 2v\ln v + \lambda v \geq 0.$$

Using $\eta^2 v$ as a test function here, we deduce

$$\lambda\int(\eta v)^2 dg \geq -4\int(\Delta v)\eta^2 v dg + \int R(\eta v)^2 dg - 2\int(\eta v)^2 \ln v dg.$$

By direct calculation

$$-4\int(\Delta v)\eta^2 v dg = 4\int|\nabla(\eta v)|^2 dg - 4\int v^2|\nabla\eta|^2 dg.$$

Hence

$$\lambda\int(\eta v)^2 dg \ge 4\int|\nabla(\eta v)|^2 dg - 4\int v^2|\nabla\eta|^2 dg + \int R(\eta v)^2 dg - 2\int(\eta v)^2\ln v dg. \tag{2.17}$$

Comparing (2.17) with (2.16) and noting that $\|\eta v\|_2 < 1$, we obtain

$$\lambda(D)\|\eta v\|_2^2 \le \lambda\|\eta v\|_2^2 + 4\int|\nabla\eta|^2 v^2 dg - \int(\eta v)^2\ln\eta^2 dg.$$

□

The next lemma is a stability result for the infimum of the Log Sobolev functional under C^2 perturbation of the metric. We believe it should be known. However, since we can not find it in the literature, we present it here.

Lemma 2.5. *Let $D \subset \mathbf{M}$ be a compact domain. For any $\epsilon > 0$, there exists $\delta > 0$ such that the following statement is true.*

Let g_1 and g_2 be two metrics on $\mathbf{M}$ such that

$$\|g_1 - g_2\|_{C^2(D,g_1)} < \delta.$$

Here $\|\cdot\|_{C^2(D,g_1)}$ stands for the C^2 norm for $(2,0)$ tensor fields under the metric g_1, restricted to the domain D. Then

$$|\lambda(D,g_1) - \lambda(D,g_2)| < \epsilon.$$

Here, for $i = 1, 2$,

$$\lambda(D,g_i) = \inf\{\int_D(4|\nabla_{g_i}v|^2 + R_{g_i}v^2 - v^2\ln v^2)dg_i \quad | \quad v \in C_0^\infty(D), \quad \|v\|_{L^2(D,g_i)=1}\}.$$

Proof.

By definition of $\lambda(D,g_1)$, there exists a function $v \in C_0^\infty(D)$ such that $\|v\|_{L^2(D,g_1)} = 1$ and that

$$\lambda(D,g_1) + \epsilon > \int_D(4|\nabla_{g_1}v|^2 + R_{g_1}v^2 - v^2\ln v^2)dg_1.$$

Recall, in local coordinate patch U with coordinate $\{x^1, ..., x^n\}$,

$$|\nabla_{g_1}v|^2 = g_1^{ij}\partial_i v\partial_j v.$$

Hence, in each local coordinate patch,

$$-\epsilon < |\nabla_{g_1}v|^2 - |\nabla_{g_2}v|^2 < \epsilon; \qquad |R_{g_1} - R_{g_2}| < \epsilon; \qquad |dg_1 - dg_2| < \epsilon$$

when $\|g_1 - g_2\|_{C^2(D,g_1)} < \delta$ with δ being sufficiently small. Since D is compact, it can be covered by finitely many local charts. Therefore, there exists $C > 0$ such that

$$\lambda(D,g_1) + \epsilon > \int_D(4|\nabla_{g_2}v|^2 + R_{g_2}v^2 - v^2\ln v^2)dg_2 - C\epsilon.$$

Consider the function $\tilde{v} = v/\|v\|_{L^2(D,g_2)}$. Then the above inequality becomes

$$\lambda(D,g_1)+\epsilon > \int_D(4|\nabla_{g_2}\tilde{v}|^2 + R_{g_2}\tilde{v}^2 - \tilde{v}^2\ln\tilde{v}^2)dg_2\,\|v\|^2_{L^2(D,g_2)} - \|v\|^2_{L^2(D,g_2)}\ln\|v\|^2_{L^2(D,g_2)} - C\epsilon.$$

Since $\|\tilde{v}\|_{L^2(D,g_2)} = 1$, we deduce

$$\lambda(D, g_1) + \epsilon > \lambda(D, g_2)\, \|v\|^2_{L^2(D,g_2)} - \|v\|^2_{L^2(D,g_2)} \ln \|v\|^2_{L^2(D,g_2)} - C\epsilon.$$

Notice that $\|v\|^2_{L^2(D,g_1)} = 1$ and $\|g_1 - g_2\|_{C^2(D,g_1)} < \delta$. Thus $|1 - \|v\|^2_{L^2(D,g_2)}| < \epsilon$ when δ is sufficiently small. Hence there exists $C > 0$ such that

$$\lambda(D, g_1) + C\epsilon > \lambda(D, g_2).$$

In the same manner, we obtain

$$\lambda(D, g_2) + C\epsilon > \lambda(D, g_1)$$

which shows

$$|\lambda(D, g_1) - \lambda(D, g_2)| < C\epsilon.$$

□

Now we are ready to give the

Proof of Theorem 1.1 (a), the existence part.

We assume $\lambda < \lambda_\infty$. First we prove that λ is finite. Since $\mathbf{M}$ has bounded geometry, it is well known (c.f. [Au], [Heb], [HV]) that the following Sobolev inequality holds: there exist positive constants S_0 depending on α, β, n such that, for all $v \in C_0^\infty(\mathbf{M})$,

$$S_0 \left(\int v^{2n/(n-2)} dg \right)^{(n-2)/n} \leq \int |\nabla v|^2 dg + \int v^2 dg.$$

Under the assumption $\|v\|_{L^2(\mathbf{M})} = 1$, a quick application of Jensen's inequality on the Sobolev inequality shows, for a constant $C = C(n, S_0)$ and all $\epsilon > 0$,

$$\int v^2 \ln v^2 dg \leq \epsilon^2 \int |\nabla v|^2 dg - \frac{n}{2} \ln \epsilon^2 + \epsilon^2 + C.$$

Taking $\epsilon = 2$ and using the assumption that the scalar curvature R is bounded, we deduce

$$\lambda = \inf\{\int_{\mathbf{M}} (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg \,|\, v \in C_0^\infty(\mathbf{M}),\ \|v\|_{L^2(\mathbf{M})} = 1\} > -\infty, \tag{2.18}$$

i.e. λ is finite.

For positive integers k, consider the domains

$$D(0, k) = \{x \in \mathbf{M} \,|\, L(x) < k\}$$

where $L = L(x)$ is the smooth function defined by (2.13), which is comparable to $d(0, x)$ when it is large. By properties of $L = L(x)$, ∂D is a C^2 boundary. Given a positive integer k, let λ_k be the best Log Sobolev constant of the ball $D(0, k)$, i.e.

$$\lambda_k = \lambda(D(0, k)) = \inf\{\int (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg \ |\ v \in C_0^\infty(D(0, k)),\ \|v\|_2 = 1\}.$$

According to [Rot], λ_k is finite and there exists a smooth extremal function v_k on $D(0, k)$, which satisfies

$$\begin{cases} 4\Delta v_k - Rv_k + 2v_k \ln v_k + \lambda_k v_k = 0, & \text{in} \quad D(0, k) \\ v_k = 0, \quad \text{on} \quad \partial D(0, k). \end{cases}$$

We mention that v_k is uniformly bounded in $C^\alpha(\mathbf{M})$ norm, i.e., there exists a positive constant C such that

$$\|v_k\|_{C^\alpha(D(0,k))} \le C. \tag{2.19}$$

A proof goes as follows. We extend v_k to a function on the whole manifold $\mathbf{M}$ by setting $v_k(x) = 0$ when $x \in \mathbf{M} - D(0,k)$. The extended function is still denoted by v_k. Then $v_k \in W^{1,2}(\mathbf{M})$, and v_k satisfies the following inequality in the weak sense

$$4\Delta v_k - R v_k + 2 v_k \ln v_k + \lambda_k v_k \ge 0, \qquad \text{in} \qquad \mathbf{M}.$$

i.e., for any nonnegative, compactly supported test function ψ, we have

$$\lambda_k \int_{\mathbf{M}} v_k \psi dg \ge \int_{\mathbf{M}} (4\nabla v_k \nabla \psi + R v_k \psi - 2\psi v_k \ln v_k) dg.$$

By Lemma 2.1, the norm $\|v_k\|_{L^\infty(\mathbf{M})}$ is uniformly bounded. Hence the original v_k in $D(0,k)$ is actually a bounded weak solution to the Poisson equation

$$\begin{cases} \Delta v_k(x) = f_k(x), \qquad x \in D(0,k) \\ v_k(x) = 0, \quad x \in \partial D(0,k) \end{cases}$$

with $\|f_k\|_{L^\infty(\mathbf{M})} \le C$. Note that $\partial D(0,k)$ is given by $L(x) = k$ and $|\nabla L(x)| + |\nabla^2 L(x)| \le C$ when $L(x)$ is large. Thus $\partial D(0,k)$ is C^2 boundary which can be expressed by a uniform C^2 function locally in geodesic balls of radius less than the injectivity radius of $\mathbf{M}$. Hence the standard elliptic theory shows (2.19) is true.

By (2.18), $\lambda_k \ge \lambda > -\infty$ and $\{\lambda_k\}$ is a decreasing sequence. Hence $\{\lambda_k\}$ is uniformly bounded by a number, say Λ. According to Lemma 2.2, there exists a point $x_k \in D(0,k)$ and a uniform constant $C = C(n, \alpha, \beta, \Lambda) > 0$ such that

$$v_k(x_k) \ge C > 0, \qquad k = 1, 2, ... \tag{2.20}$$

We consider 2 cases.

Case 1. $\{x_k\}$ is a bounded sequence in $\mathbf{M}$, i.e. $d(x_k, 0)$ is uniformly bounded.

By Lemma 2.1, the sequence $\{v_k\}$ of extended functions is uniformly bounded in L^∞ norm, $k = 2, 3, ...$ By (2.19) we can find a subsequence, still denoted by $\{v_k\}$, which converges in C^α_{loc} norm to a smooth, nonnegative function $v \in C^\infty(\mathbf{M})$ that solves the equation

$$4\Delta v - Rv + 2v \ln v + \lambda v = 0.$$

The lower bound in (2.20) ensures that v is a positive solution. Moreover $\|v\|_{L^2(\mathbf{M})} \le 1$ by Fatou's Lemma. By Lemma 2.3, there exist positive constants a and A such that

$$v(x) \le A e^{-a d^2(x,0)} \qquad x \in \mathbf{M}.$$

The classical volume comparison theorem tells us that $|B(0,k)|_g$ grows at most like e^{ck}, where c depends on the curvature bound α and n. Hence we can multiply the above equation by v and perform integration by parts to deduce

$$L(v,g) = \int_{\mathbf{M}} [4|\nabla v|^2 + R v^2 - v^2 \ln v^2] dg = \lambda \int_{\mathbf{M}} v^2 dg. \tag{2.21}$$

If $\int_{\mathbf{M}} v^2 dg = 1$, then v is an extremal function of the Log Sobolev functional L, and the proof Theorem 1.1 (a) is done. So we suppose $\int_{\mathbf{M}} v^2 dg < 1$. We consider the function

$$\tilde{v} = \frac{v}{\|v\|_{L^2(\mathbf{M})}}.$$

Then $\|\tilde{v}\|_{L^2(\mathbf{M})} = 1$ and (2.21) infers

$$\begin{aligned}
\lambda = L(v,g)\|v\|^{-2}_{L^2(\mathbf{M})} &= \frac{\int_{\mathbf{M}}[4|\nabla v|^2 + Rv^2 - v^2\ln v^2]dg}{\|v\|^2_{L^2(\mathbf{M})}} \\
&= \int_{\mathbf{M}}[4|\nabla \tilde{v}|^2 + R\tilde{v}^2 - \tilde{v}^2\ln \tilde{v}^2]dg - \ln\|v\|^2_{L^2(\mathbf{M})} \\
&\geq \lambda - \ln\|v\|^2_{L^2(\mathbf{M})}.
\end{aligned}$$

The last step is due to the definition that λ is the infimum of the Log Sobolev functional. If the assumption $\int_{\mathbf{M}} v^2 dg < 1$ is valid, we would get the contradiction $\lambda > \lambda$. Hence $\int_{\mathbf{M}} v^2 dg = 1$ and v is indeed an extremal. This finishes the proof in Case 1.

Case 2. $\{x_k\}$ is an unbounded sequence in $\mathbf{M}$.

Since $\mathbf{M}$ has bounded geometry, by Hamilton's compactness theorem, the pointed manifolds $(\mathbf{M}, x_k, g)$ converges in C^∞_{loc} topology (also called Cheeger-Gromov sense), to a complete limit manifold $(M_\infty, x_\infty, g_\infty)$. This limit manifold also has bounded geometry.

Recall $v_k(\geq 0)$ solves

$$\begin{cases} 4\Delta v_k - Rv_k + 2v_k\ln v_k + \lambda_k v_k = 0, \qquad \text{in} \qquad D(0,k) \\ v_k = 0, \qquad \text{on} \qquad \partial D(0,k). \end{cases}$$

We extend v_k to a function on the whole manifold $\mathbf{M}$ by setting $v_k(x) = 0$ when $x \in \mathbf{M} - D(0,k)$. The extended function is still denoted by v_k. Then, as in Case 1, $v_k \in C^\alpha(\mathbf{M}) \cap W^{1,2}(\mathbf{M})$, and v_k satisfies the following inequality in the weak sense

$$4\Delta v_k - Rv_k + 2v_k\ln v_k + \lambda_k v_k \geq 0, \qquad \text{in} \qquad \mathbf{M}.$$

Since v_k is nonnegative and uniformly bounded by Lemma 2.1, the standard elliptic theory shows that a subsequence of $\{v_k\}$, converges in C^α_{loc} sense to a function $v \in C^\alpha(\mathbf{M}_\infty) \cap W^{1,2}(\mathbf{M}_\infty)$. Moreover v satisfies the following inequality in the weak sense

$$4\Delta v - Rv + 2v\ln v + \lambda v \geq 0, \qquad \text{in} \qquad \mathbf{M}_\infty.$$

i.e., for any nonnegative, compactly supported test function ψ, we have

$$\lambda \int_{\mathbf{M}_\infty} v\psi dg_\infty \geq \int_{\mathbf{M}_\infty} (4\nabla v \nabla \psi + Rv\psi - 2\psi v\ln v) dg_\infty.$$

Here the Laplacian Δ, the gradient ∇ and the scalar curvature R are with respect to the limiting metric g_∞. Since $v_k(x_k)$ converges to $v(x_\infty)$, by (2.20), we also know that

$$v(x_\infty) > C > 0. \tag{2.22}$$

By Lemma 2.3 and Fatou Lemma, there hold the bounds

$$v(x) \leq Ae^{-ad^2(x,x_\infty,g_\infty)}, \quad x \in \mathbf{M}_\infty; \qquad \int_{\mathbf{M}_\infty} v^2(x)dg_\infty \leq 1. \tag{2.23}$$

Let $r > 0$ be a large number to be fixed later. Define, on the manifold (M_∞, g_∞) and under the metric g_∞,

$$\lambda(B(x_\infty, r)) = \inf\{\int (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg_\infty \mid v \in C_0^\infty(B(x_\infty, r)),\ \|v\|_2 = 1\}.$$

We choose a smooth cut-off function $\eta \in C_0^\infty(B(x_\infty, r))$ such that $0 \le \eta \le 1$, $\eta = 1$ on $B(x_\infty, r/2)$ and that $|\nabla \eta| \le C/r$. By Lemma 2.4, it holds

$$\lambda(B(x_\infty, r)) \le \lambda + 4\frac{\int v^2|\nabla\eta|^2 dg_\infty}{\int (v\eta)^2 dg_\infty} - \frac{\int (v\eta)^2 \ln \eta^2 dg_\infty}{\int (v\eta)^2 dg_\infty}. \tag{2.24}$$

By (2.22) and the fact that v is in $C^\alpha(M_\infty)$, we can find a positive constant $c > 0$ such that

$$\int (v\eta)^2 dg_\infty \ge \int_{B(x_\infty, r/2)} v^2 dg_\infty \ge c.$$

From this and (2.24), using properties of η, we deduce

$$\lambda(B(x_\infty, r)) \le \lambda + C(1 + 1/r) \int_{B(x_\infty, r) - B(x_\infty, r/2)} v^2 dg_\infty.$$

By (2.23) and the classical volume comparison theorem, this implies

$$\lambda(B(x_\infty, r)) \le \lambda + C(1 + 1/r) e^{-ar^2/4} e^{c\alpha r}.$$

Here, as before α is the bound on the curvature tensor. Thus, for any $\epsilon > 0$, there exists $r_0 > 0$ such that

$$\lambda = \lambda(\mathbf{M}) \ge \lambda(B(x_\infty, r)) - \epsilon \tag{2.25}$$

when $r \ge r_0$.

By definition of $(M_\infty, x_\infty, g_\infty)$ as a limit manifold, for any $\delta > 0$, when k is sufficiently large, there exists a diffeomorphism F from $B(x_\infty, r)$ onto an open set $U \subset \mathbf{M}$, which contains x_k, such that $(F^*)^{-1} g_\infty$ and g are δ close in C^∞ topology, when they are restricted to U. By Lemma 2.5, we have, when δ is sufficiently small,

$$\lambda(B(x_\infty, r)) = \lambda(B(x_\infty, r), g_\infty) = \lambda(U, (F^*)^{-1} g_\infty) > \lambda(U, g) - \epsilon. \tag{2.26}$$

By definition of U, we know that for any $x \in U$,

$$d(x, x_k, (F^*)^{-1} g_\infty) < r$$

which implies, since $(F^*)^{-1} g_\infty$ and g are δ close,

$$d(x, x_k, g) < (1 + C\sqrt{\delta}) r.$$

Hence, when δ is sufficiently small, it holds

$$U \subset B(x_k, 2r, g).$$

This and (2.26) tell us that

$$\lambda(B(x_\infty, r)) > \lambda(B(x_k, 2r, g), g) - \epsilon.$$

Recall that $d(x_k, 0, g) \to \infty$ when $k \to \infty$. Therefore, when k is large,

$$B(x_k, 2r, g) \subset \mathbf{M} - B(0, d(x_k, 0, g)/2, g).$$

By definition of λ_∞, we know that

$$\lambda(B(x_k, 2r, g), g) > \lambda_\infty - \epsilon$$

when k is sufficiently large. So we get

$$\lambda(B(x_\infty, r)) > \lambda_\infty - 2\epsilon.$$

By (2.25), we finally deduce

$$\lambda = \lambda(\mathbf{M}) > \lambda_\infty - 3\epsilon.$$

Since ϵ can be sufficiently small, we have reached a contradiction with the assumption that $\lambda < \lambda_\infty$. This shows that Case 2 can not happen, and only Case 1 occurs, implying that an extremal exists.

The bound for the extremal v in the theorem is already proven in Lemma 2.3. This proves part (a) of the theorem. □

3. Proof of the theorem 1.1 (b), the nonexistence part

The proof is done by constructing a concrete 3 manifold on which the Log Sobolev functional does not have an extremal. In order to present the main idea of the construction, we informally describe a crude example of a disconnected manifold of such kind.

Example 3.1. Let (M_k, g_k), $k = 1, 2, ...$, be a sequence of compact manifolds without boundary and let λ_k be the infimum of the Log Sobolev functional on M_k. We assume that λ_k is a strictly decreasing sequence bounded from below by a finite number. For instance we can take $M_k = (1 + k^{-2})(S^1 \times S^1)$, the flat 2 torus whose metric is the standard one scaled by the factor $1 + k^{-2}$. Let M be the disjoint union of M_k. We now prove that the Log Sobolev functional does not have an extremal on M. Suppose for contradiction that v is an extremal of the Log Sobolev functional on M, whose infimum is λ. Then $\lambda < \lambda_k$ and

$$\lambda = L(v, g) = \Sigma_{k=1}^{\infty} \int_{M_k} (4|\nabla v|^2 + R_k v^2 - v^2 \ln v^2) dg_k. \tag{3.1}$$

Here R_k is the scalar curvature of (M_k, g_k). Without loss of generality, we can assume that $v|_{M_k}$ is not identically zero for $k = 1, 2, 3, ...$ Otherwise, we just delete those M_k where $v|_{M_k}$ is identically zero. Write

$$v_k = \frac{v|_{M_k}}{\|v|_{M_k}\|_{L^2(M_k, g_k)}}.$$

Then, $\|v_k\|^2_{L^2(M_k, g_k)} = 1$ and

$$\begin{aligned}
&\int_{M_k} (4|\nabla v|^2 + R_k v^2 - v^2 \ln v^2) dg_k \\
&= \|v|_{M_k}\|^2_{L^2(M_k, g_k)} \int_{M_k} (4|\nabla v_k|^2 + R_k v_k^2 - v_k^2 \ln v_k^2) dg_k - \|v|_{M_k}\|^2_{L^2(M_k, g_k)} \ln \|v|_{M_k}\|^2_{L^2(M_k, g_k)} \\
&\geq \|v|_{M_k}\|^2_{L^2(M_k, g_k)} \int_{M_k} (4|\nabla v_k|^2 + R_k v_k^2 - v_k^2 \ln v_k^2) dg_k.
\end{aligned}$$

Here we used the fact that $\|v|_{M_k}\|^2_{L^2(M_k,g_k)} \le \|v\|^2_{L^2(M)} = 1$. Hence

$$\int_{M_k} (4|\nabla v|^2 + R_k v^2 - v^2 \ln v^2) dg_k \ge \|v|_{M_k}\|^2_{L^2(M_k,g_k)} \lambda_k.$$

Substituting this to (3.1), we deduce

$$\lambda \ge \Sigma^\infty_{k=1} \|v|_{M_k}\|^2_{L^2(M_k,g_k)} \lambda_k.$$

Notice that

$$1 = \|v\|^2_{L^2(M)} = \Sigma^\infty_{k=1} \|v|_{M_k}\|^2_{L^2(M_k,g_k)}.$$

Multiplying this equality by λ and subtracting the last inequality, we find that

$$\Sigma^\infty_{k=1} \|v|_{M_k}\|^2_{L^2(M_k,g_k)} (\lambda_k - \lambda) \le 0,$$

which is a contradiction with the fact that $\lambda_k > \lambda$. Hence no such extremal v exists.

The manifold M in this example is disconnected and therefore it can not serve as a proof of the theorem. However, building on the main idea from this example, we will construct a manifold $\mathbf{M}$ which is a connected sum of infinitely many copies of compact manifolds, each of which can be graphically described as a ball with a handle or just a "hand bag". See the figure in Step 4 of the proof. The basic components of the manifold are: round necks, truncated S^3, and tubes whose cross sections are the flat torus $S^1 \times S^1$. By studying the behavior of the Log Sobolev functional when these components are pasted together, we will eventually show that the Log Sobolev functional does not have an extremal.

First let us introduce some notations.

Definition 3.1. *(Round necks and flat tubes)*

Let h, A, B *be real numbers, we use* $N = N(h, A, B)$ *to denote the round neck* $h^2 S^2 \times [A, B]$ *with the product metric* $g = h^2 g_{S^2} \times g_{R^1}$. *Here* g_{S^2} *is the standard round metric on* S^2 *with radius* 1; g_{R^1} *is the Euclidean metric on* R^1; *and* h^2 *scales* g_{S^2} *only. For convenience, we also normalize the scalar curvature corresponding to* g_{S^2} *to be* 1. *Let* $x \in N(h, A, B)$. *We use* $x = (x_1, x_2, x_3)$ *as a coordinate for* x, *where* $(x_1, x_2) \in S^2$ *and* $x_3 \in [A, B]$.

If $A = 0$, *we will use* $N(h, B)$ *to denote* $N(h, A, B)$.

We use $H = H(h, A, B)$ *to denote the flat tube* $h^2(S^1 \times S^1) \times [A, B]$ *with the product metric* $g = h^2 g_{S^1\times S^1} \times g_{R^1}$. *Here* $g_{S^1\times S^1}$ *is the standard flat metric on* $S^1 \times S^1$ *so that the radius of* S^1 *is* 1; g_{R^1} *is the Euclidean metric on* R^1; *and* h^2 *scales* $g_{S^1\times S^1}$ *only. Let* $x \in H(h, A, B)$. *We use* $x = (x_1, x_2, x_3)$ *as a coordinate for* x, *where* $(x_1, x_2) \in S^1 \times S^1$ *and* $x_3 \in [A, B]$.

We need a number of lemmas again.

Lemma 3.1. *Let* v *be a bounded, positive subsolution to the equation (2.1) in the round neck* $N = h^2 S^2 \times [-l, l]$. *i.e.*

$$4\Delta v - Rv + 2v \ln v + \lambda v \ge 0.$$

Suppose $\lambda \le 0$, $h \in (0, 1]$, $l \ge 2$ *and that* $\|v\|_{L^2(N)} \le 1$. *Then there exists a positive constant* C *which is independent of* h *such that*

$$v^2(x) \le C \int_{B(x,1)} v^2 dg$$

when $x \in h^2S^2 \times [-l+1, l-1]$.

Proof.

The result in this lemma and the proof are analogous to that in Lemma 2.1. However, there is difference, namely the constant C in the lemma is independent of $h \in (0,1]$.

First, we claim that there exists a positive constant S_0, independent of h, such that such that,

$$S_0 \left(\int u^{2n/(n-2)} dg \right)^{(n-2)/n} \le \int (4|\nabla u|^2 + Ru^2) dg, \qquad n = 3, \tag{3.2}$$

for all $u \in C_0^\infty(h^2S^2 \times [-l,,l])$. Here is a quick proof of the claim. Consider the infinite round neck $S^2 \times h^{-2}R^1$. Here $h^{-2}R^1$ is R^1equipped with the scaled metric $h^{-2}g_{R^1}$. Note the curvature bounds and the lower bound of injectivity radius are independent of h. i.e. the necks have uniformly bounded geometry. By [Au], there exists a positive constant S_0 such that

$$S_0 \left(\int u^{2n/(n-2)} dg \right)^{(n-2)/n} \le \int (|\nabla u|^2 + u^2) dg$$

for all $u \in C_0^\infty(S^2 \times h^{-2}R^1)$. Notice that the scalar curvature of $S^2 \times h^{-2}R^1$ is the constant 1. Hence

$$S_0 \left(\int u^{2n/(n-2)} dg \right)^{(n-2)/n} \le \int (4|\nabla u|^2 + Ru^2) dg$$

for all $u \in C_0^\infty(S^2 \times h^{-2}R^1)$. But this Sobolev inequality is scaling invariant. Hence, for all $u \in C_0^\infty(h^2S^2 \times R^1)$, inequality (3.2) holds, proving the claim.

Since v is a subsolution of (2.1) and $\lambda \le 0$ by assumption, given any $p \ge 1$, it is easy to see that

$$-4\Delta v^p + pRv^p \le 2pv^p \ln v.$$

We select a smooth cut off function ϕ supported in $h^2S^2 \times [-l,,l]$. Writing $w = v^p$ and using $w\phi^2$ as a test function in the above inequality, we deduce

$$4 \int \nabla(w\phi^2)\nabla w dg + p \int R(w\phi)^2 dg \le 2p \int (w\phi)^2 \ln v dg.$$

Since the scalar curvature is positive, this shows

$$4 \int \nabla(w\phi^2)\nabla w dg + \int R(w\phi)^2 dg \le 2p \int (w\phi)^2 \ln v^2 dg,$$

which induces, after integration by parts,

$$\int (4|\nabla(w\phi)|^2 + R(w\phi)^2) dg \le 4 \int |\nabla \phi|^2 w^2 dg + 2p \int (w\phi)^2 \ln v^2 dg.$$

Applying (3.2) on the left hand side, we deduce

$$S_0 \left(\int (w\phi)^{2n/(n-2)} dg \right)^{(n-2)/n} \le 4 \int |\nabla \phi|^2 w^2 dg + 2p \int (w\phi)^2 \ln v^2 dg.$$

Now pick $x \in h^2S^2 \times [-l+1, l-1]$. Then $B(x,1) \subset h^2S^2 \times [-l+1, l-1]$. Now we choose ϕ as suitable cut-off functions supported in $B(x,1)$. The rest of the proof of the lemma is the same as the proof of Lemma 2.1 after (2.6), with λ there taken as 0. $\square$

The next lemma says that if v is a solution of (2.1) in a very long round neck, whose L^2 norm is less than 1, then v is exponentially small in the middle section of the neck.

Lemma 3.2. *There exists $h_0 \in (0,1]$ such that the following statement holds for all $h \in (0, h_0]$. Let v be a smooth positive solution to the equation (2.1) in the round neck $N = h^2 S^2 \times [-l, l]$. Suppose $\lambda \le 0$, $l \ge 2$ and that $\|v\|_{L^2(N)} \le 1$. Then there exist positive constants a and A, independent of h, such that*

$$\int_{h^2S^2\times[-l/2,l/2]} v^2 dg \le Ae^{-al}\,[\int_{h^2S^2\times[-l,-l+2]} v^2 dg + \int_{h^2S^2\times[l-2,l]} v^2 dg]$$

and

$$v(x) \le Ae^{-al}, \quad x \in h^2S^2 \times [-l/2, l/2].$$

Proof.

By the previous lemma, for $x \in h^2S^2 \times [-l+1, l-1]$, we have a constant C such that

$$v(x) \le C.$$

Note the scalar curvature $R = 1/h^2$. Hence there exists $h_0 \in (0,1]$ such that if $h \in (0, h_0]$ then

$$R/2 - 2\ln v \ge 1/(2h_0^2) - 2\ln C \ge 0.$$

Combining this with equation (2.1) i.e.

$$4\Delta v - Rv + 2v\ln v + \lambda v = 0,$$

we find that v satisfies the inequality

$$\Delta v - \frac{1}{8h_0^2} v \ge 0 \quad \text{in} \quad h^2S^2 \times [-l+1, l-1]. \tag{3.3}$$

Here we have used the assumption that $\lambda \le 0$.

We pick a cut off function $\phi \in C_0^\infty(N)$, satisfying the following requirements.

$$\phi(x) = \phi(x_1, x_2, x_3) = \begin{cases} 0, & x_3 \in [-l, -l+1] \cup [l-1, l], \\ \text{a number in} \quad (0,1), & x_3 \in [-l+1, -l+2] \cup [l-2, l-1] \\ 1, & x_3 \in [-l+2, l-2]. \end{cases}$$

We also require that $|\nabla\phi| \le 4$. Here we recall that x_3 is the longitudinal component of the coordinate of the point x in the neck N, as described in Definition 3.1. See the figure below.



Let a be a positive number to be determined later. Using $e^{a(l-|x_3|)}\phi^2 v$ as a test function on (3.3) and performing integration by parts, we find that

(3.4)

$$\begin{aligned}&\frac{1}{8h_0^2}\int e^{a(l-|x_3|)}\phi^2 v^2 dg \le \int e^{a(l-|x_3|)}\phi^2 v\Delta v dg\\ &= -\int e^{a(l-|x_3|)}\phi^2|\nabla v|^2 dg - 2\int e^{a(l-|x_3|)} v\phi\nabla\phi\nabla v dg - \int e^{a(l-|x_3|)}\nabla(a(l-|x_3|))\nabla v v\phi^2 dg\\ &\equiv -Y_1 - Y_2 - Y_3.\end{aligned}$$

We need to bound $|Y_2|$ and $|Y_3|$.

First we notice

$$\begin{aligned}|Y_2| &\le 2\int e^{a(l-|x_3|)} v\phi|\nabla\phi\nabla v| dg\\ &\le \frac{1}{4}\int e^{a(l-|x_3|)}\phi^2|\nabla v|^2 dg + 4\int e^{a(l-|x_3|)} v^2|\nabla\phi|^2 dg.\end{aligned}$$

Therefore

$$|Y_2| \le \frac{1}{4}Y_1 + 4\int_{supp\nabla\phi} e^{a(l-|x_3|)} v^2 dg. \tag{3.5}$$

Next

$$\begin{aligned}|Y_3| &\le a\int e^{a(l-|x_3|)}|\nabla v| v\phi^2 dg\\ &\le \frac{a}{2}\int e^{a(l-|x_3|)}\phi^2 v^2 dg + \frac{a}{2}\int e^{a(l-|x_3|)}|\nabla v|^2\phi^2 dg\\ &= \frac{a}{2}\int e^{a(l-|x_3|)}\phi^2 v^2 dg + \frac{a}{2}Y_1.\end{aligned}$$

Choosing $a \le 1$ and substituting this and (3.5) into (3.4), we deduce

$$\frac{1}{8h_0^2}\int e^{a(l-|x_3|)}\phi^2 v^2 dg \le 4\int_{supp\nabla\phi} e^{a(l-|x_3|)} v^2 dg + \frac{a}{2}\int e^{a(l-|x_3|)}\phi^2 v^2 dg.$$

Taking $a = \min\{1, \frac{1}{8h_0^2}\}$, we arrive at

$$\int e^{a(l-|x_3|)}\phi^2 v^2 dg \le C h_0^2 \int_{supp\nabla\phi} e^{a(l-|x_3|)} v^2 dg. \tag{3.6}$$

Observe that when $x \in supp\nabla\phi$ we have

$$0 \le l - |x_3| \le 2.$$

Also, when $x \in h^2 S^2 \times [-2l/3, 2l/3]$, i.e. when $-2l/3 \le x_3 \le 2l/3$, we have

$$l - |x_3| \ge l/3, \qquad \phi(x) = 1.$$

Therefore (3.6) implies

$$\int_{h^2 S^2\times[-2l/3,2l/3]} v^2 dg \le C h_0^2 e^{2a} e^{-al/3}\,[\int_{h^2 S^2\times[-l,-l+2]} v^2 dg + \int_{h^2 S^2\times[l-2,l]} v^2 dg]$$

which yields the desired integral bound, after adjusting the coefficients. The pointwise bound in the lemma is an immediate consequence the integral bound and Lemma 3.1 □

Let v again be a positive solution of (2.1) in a very long round neck, whose L^2 norm is less than 1. The next lemma says that if v vanishes at one end of the neck, then v is exponentially small near that end.

Lemma 3.3. *There exists $h_0 \in (0,1]$ such that the following statement holds for all $h \in (0, h_0]$. Let v be a smooth positive solution to the equation (2.1) in the round neck $N = h^2S^2 \times [0,l]$. Suppose $\lambda \le 0$, $l \ge 2$ and that $\|v\|_{L^2(N)} \le 1$. Suppose also $v(x) = 0$ when $x \in h^2S^2 \times \{l\}$. i.e. v vanishes at the right end of the neck. Then there exist positive constants a and A, independent of h, such that*

$$\int_{h^2S^2\times[l/2,l]} v^2 dg \le Ae^{-al} \int_{h^2S^2\times[0,1]} v^2 dg.$$

Proof.

We extend $v = v(x)$ to a function on the longer neck $h^2S^2 \times [0, l+1]$ by assigning $v(x) = 0$ when $x_3 \ge l$. Since $v(x) = 0$ when $x_3 = l$, it is easy to see that the extended v is a subsolution to (2.1) on $h^2S^2 \times [0, l+1]$. By Lemma 3.1, for $x \in h^2S^2 \times [1, l]$, there exists a constant C such that

$$v(x) \le C.$$

Since the scalar curvature $R = 1/h^2$, there exists $h_0 \in (0,1]$ such that if $h \in (0, h_0]$ then

$$R/2 - 2\ln v \ge 1/(2h_0^2) - 2\ln C \ge 0.$$

Combining this with equation (2.1) i.e.

$$4\Delta v - Rv + 2v\ln v + \lambda v = 0.$$

we find that v satisfies the inequality

$$\Delta v - \frac{1}{8h_0^2} v \ge 0 \quad \text{in} \quad h^2S^2 \times [1,l]. \tag{3.7}$$

Here we have again used the assumption that $\lambda \le 0$.

We pick a cut off function $\phi \in C_0^\infty(N)$, satisfying $|\nabla\phi| \le 4$ and the following requirements.

$$\phi(x) = \phi(x_1, x_2, x_3) = \begin{cases} 0, & & x_3 \in [0,1], \\ \text{a number in} & (0,1), & x_3 \in [1,2] \\ 1, & & x_3 \in [2,l]. \end{cases}$$

Let a be a positive number to be determined later. Using $e^{ax_3}\phi^2 v$ as a test function on (3.7) and performing integration by parts, we find that

$$\begin{aligned} &\frac{1}{8h_0^2}\int e^{ax_3}\phi^2 v^2 dg \le \int e^{ax_3}\phi^2 v\Delta v dg \\ &= -\int e^{ax_3}\phi^2|\nabla v|^2 dg - 2\int e^{ax_3} v\phi\nabla\phi\nabla v dg - \int e^{ax_3}\nabla(ax_3)\nabla v v\phi^2 dg \\ &\equiv -Y_1 - Y_2 - Y_3. \end{aligned} \tag{3.8}$$

Note that boundary terms vanish since $v = 0$ at the right end of the neck and $\phi = 0$ at the left end. Let us bound $|Y_2|$ and $|Y_3|$.

First we notice

$$\begin{aligned}|Y_2| &\le 2\int e^{ax_3} v\phi|\nabla\phi\nabla v|dg\\ &\le \frac{1}{4}\int e^{ax_3}\phi^2|\nabla v|^2dg + 4\int e^{ax_3}v^2|\nabla\phi|^2dg.\end{aligned}$$

Therefore

$$|Y_2| \le \frac{1}{4}Y_1 + 4\int_{supp\nabla\phi} e^{ax_3}v^2dg. \tag{3.9}$$

Next

$$\begin{aligned}|Y_3| &\le a\int e^{ax_3}|\nabla v|v\phi^2dg\\ &\le \frac{a}{2}\int e^{ax_3}\phi^2v^2dg + \frac{a}{2}\int e^{ax_3}|\nabla v|^2\phi^2dg\\ &= \frac{a}{2}\int e^{ax_3}\phi^2v^2dg + \frac{a}{2}Y_1.\end{aligned}$$

Choosing $a \le 1$ and substituting this and (3.9) into (3.8), we deduce

$$\frac{1}{8h_0^2}\int e^{ax_3}\phi^2v^2dg \le 4\int_{supp\nabla\phi} e^{ax_3}v^2dg + \frac{a}{2}\int e^{ax_3}\phi^2v^2dg.$$

Taking $a = \min\{1, \frac{1}{8h_0^2}\}$, we arrive at

$$\int e^{ax_3}\phi^2v^2dg \le Ch_0^2\int_{supp\nabla\phi} e^{ax_3}v^2dg. \tag{3.10}$$

Observe that when $x \in supp\nabla\phi$ we have

$$0 \le x_3 \le 1.$$

Also, when $x \in h^2S^2 \times [l/2, l]$, we have

$$x_3 \ge l/2, \qquad \phi(x) = 1.$$

Therefore (3.10) implies

$$\int_{h^2S^2\times[l/2,l]} v^2dg \le Ch_0^2e^ae^{-al/2}\int_{h^2S^2\times[0,1]} v^2dg,$$

proving the lemma. □

The following lemma is similar to Lemma 2.4. The difference is that we are comparing the infimum of the Log Sobolev functionals on two different domains in this lemma. The proof is almost identical.

Lemma 3.4. *Let E and F be two domains of $\mathbf{M}$ such that $E \subset F$ and that E is compact. Let $v \in W_0^{1,2}(F)$, $\|v\|_{L^2(F)} = 1$ be an extremal of $\lambda(F)$ so that it is a smooth positive solution of the equation*

$$4\Delta v - Rv + 2v\ln v + \lambda(F)v = 0.$$

For any smooth cut-off function η such that $\eta v \in C_0^\infty(E)$ and $0 \le \eta \le 1$, it holds

$$\lambda(E) \le \lambda(F) + 4\frac{\int v^2|\nabla\eta|^2dg}{\int(v\eta)^2dg} - \frac{\int(v\eta)^2\ln\eta^2dg}{\int(v\eta)^2dg}.$$

Proof.

Since $\eta v/\|\eta v\|_2 \in C_0^\infty(E)$ and its L^2 norm is 1, we have, by definition,

$$\lambda(E) \le \int \left[4\frac{|\nabla(\eta v)|^2}{\|\eta v\|_2^2} + R\frac{(\eta v)^2}{\|\eta v\|_2^2} - \frac{(\eta v)^2}{\|\eta v\|_2^2} \ln \frac{(\eta v)^2}{\|\eta v\|_2^2} \right] dg.$$

This implies

$$\lambda(E)\|\eta v\|_2^2 \le \int \left[4|\nabla(\eta v)|^2 + R(\eta v)^2 - (\eta v)^2 \ln(\eta v)^2 \right] dg + \|\eta v\|_2^2 \ln \|\eta v\|_2^2. \tag{3.11}$$

On the other hand, v is a smooth positive solution of the equation

$$4\Delta v - Rv + 2v \ln v + \lambda(F) v = 0.$$

Using $\eta^2 v$ as a test function for the equation, we find

$$\lambda(F) \int (\eta v)^2 dg = -4 \int (\Delta v) \eta^2 v dg + \int R(\eta v)^2 dg - 2 \int (\eta v)^2 \ln v dg.$$

Using integration by parts, we deduce

$$-4 \int (\Delta v) \eta^2 v dg = 4 \int |\nabla(\eta v)|^2 dg - 4 \int v^2 |\nabla \eta|^2 dg.$$

Hence

$$\lambda(F) \int (\eta v)^2 dg = 4 \int |\nabla(\eta v)|^2 dg - 4 \int v^2 |\nabla\eta|^2 dg + \int R(\eta v)^2 dg - 2 \int (\eta v)^2 \ln v dg. \tag{3.12}$$

Comparing (3.12) with (3.11) and noting that $\|\eta v\|_2 \le 1$, we obtain

$$\lambda(E)\|\eta v\|_2^2 \le \lambda(F)\|\eta v\|_2^2 + 4 \int |\nabla \eta|^2 v^2 dg - \int (\eta v)^2 \ln \eta^2 dg.$$

□

The following lemma says that if a domain E contains a round neck of length l and F is the extension of E, which is obtained by pasting a segment of the round neck with length 1, then $|\lambda(E) - \lambda(F)|$ is exponentially small.

Lemma 3.5. *Let $E \subset \mathbf{M}$ be a compact domain such that*

$$E = X_0 \cup N(h, l)$$

which is the connected, non-overlapping union of a domain X_0 with the round neck $N(h,l) = h^2 S^2 \times [0, l]$. Let

$$F = X_0 \cup N(h, l+1)$$

which is the connected, non-overlapping union of X_0 with the round neck $N(h, l+1) = h^2 S^2 \times [0, l+1]$. There is $h_0 \in [0,1]$ and $l_0 > 0$ such that for all $h \in [0, h_0]$ and $l \ge l_0$, the following statement holds:

If $\lambda(E) \le 0$, then there exist positive numbers a and A such that

$$\lambda(F) \ge \lambda(E) - Ae^{-al}.$$

Proof.

First let us see the figure depicting E and F below.



Pick a smooth cut off function η such that $|\nabla\eta| \le 4$ and that

$$\eta = \eta(x) = \begin{cases} 0, & x \in h^2S^2 \times [l, l+1] \\ \text{a number in} \quad (0,1), & x \in h^2S^2 \times [l-1, l] \\ 1, & x \in F - (h^2S^2 \times [l-1, l+1]). \end{cases}$$

Let v be an extremal for $\lambda(F)$, which exists since F is compact. Then $\eta v \in C_0^\infty(E)$. By Lemma 3.4, we have

$$\lambda(E) \le \lambda(F) + 4\frac{\int v^2|\nabla\eta|^2 dg}{\int (v\eta)^2 dg} - \frac{\int (v\eta)^2 \ln \eta^2 dg}{\int (v\eta)^2 dg}. \tag{3.13}$$

Observe that

$$\int (v\eta)^2 dg = \int v^2 dg - \int v^2(1-\eta^2) dg \ge 1 - \int_{h^2S^2\times[l-1,l+1]} v^2 dg.$$

Using Lemma 3.3 on $h^2S^2 \times [0, l+1]$, we infer, for some positive numbers a and A, that

$$\int_{h^2S^2\times[l-1,l+1]} v^2 dg \le Ae^{-al}.$$

Hence

$$\int (v\eta)^2 dg \ge 1 - Ae^{-al}.$$

Also notice that

$$\int v^2|\nabla\eta|^2 dg \le 16 \int_{h^2S^2\times[l-1,l]} v^2 dg \le 16Ae^{-al},$$

and

$$|\int (v\eta)^2 \ln \eta^2 dg| \le e^{-1} \int_{h^2S^2\times[l-1,l]} v^2 dg \le Ae^{-al}.$$

Substituting the last three inequalities into (3.13), we deduce

$$\lambda(E) \le \lambda(F) + C\frac{Ae^{-al}}{1 - Ae^{-al}}.$$

Therefore, there exists $l_0 > 0$ such that for all $l \ge l_0$, we have

$$\lambda(E) \le \lambda(F) + Ae^{-al}$$

for some constant $A > 0$, whose value may have been adjusted from the last line. □

The following lemma says that the infimum of the Log Sobolev functional on a flat tube goes to $-\infty$ when the cross section of the tube goes to 0.

Lemma 3.6. *Let $H = H(h, 0, 1) = h^2(S^1 \times S^1) \times [0, 1]$ be a flat tube given in Definition 3.1. Then $\lambda(H(h, 0, 1)) \to -\infty$ when $h \to 0$.*

Proof.

Given $x \in H(h, 0, 1)$, let (x_1, x_2, x_3) be its coordinate described in Definition 3.1. Consider the one variable function

$$v = v(x_3) = \begin{cases} 4\frac{\sqrt{3}}{\sqrt{8}\pi h}x_3, & x_3 \in [0, 1/4], \\ \frac{\sqrt{3}}{\sqrt{8}\pi h}, & x_3 \in [1/4, 3/4], \\ \frac{\sqrt{3}}{\sqrt{8}\pi h}[1 - 4(x_3 - 3/4)], & x_3 \in [3/4, 1]. \end{cases}$$

We compute

$$\int_{H(h,0,1)} v^2 dg = 4\pi^2 h^2 \int_0^1 v^2 dx_3 = \frac{3}{8\pi^2 h^2} 4\pi^2 h^2 (2\int_0^{1/4} 16x_3^2 dx_3 + \frac{1}{2}) = 1,$$

$$\int_{H(h,0,1)} |\nabla v|^2 dg = 4\pi^2 h^2 \int_0^1 |\partial_{x_3} v|^2 dx_3 = \frac{3}{8\pi^2 h^2} 4\pi^2 h^2 (2\int_0^{1/4} 16 dx_3) = 12,$$

$$\begin{aligned} \int_{H(h,0,1)} v^2 \ln v^2 dg &= 4\pi^2 h^2 \int_0^1 v^2 \ln v^2 dx_3 \\ &= \frac{3}{8\pi^2 h^2} 4\pi^2 h^2 [2\int_0^{1/4} 16x_3^2 \ln(\frac{3}{8\pi^2 h^2} 16x_3^2) dx_3 + \int_{1/4}^{3/4} \ln(\frac{3}{8\pi^2 h^2}) dx_3] \\ &= -\frac{3}{4} \ln h^2 + c \end{aligned}$$

where c is a constant independent of h.

Since the scalar curvature is zero, these computation imply

$$\lambda(H(h, 0, 1)) \le \int_{H(h,0,1)} (4|\nabla v|^2 - v^2 \ln v^2) dg = \frac{3}{4} \ln h^2 + c.$$

This shows $\lambda(H(h, 0, 1)) \to -\infty$ when $h \to 0$. □

Now we are ready to give

Proof of Theorem 1.1 (b).

As mentioned earlier we will construct a noncompact manifold with bounded geometry such that the Log Sobolev functional does not have an extremal. The manifold is a connected sum of infinitely many components connected by increasingly long round necks. Each of the component shapes like a hand bag. The handle of a hand bag is a flat tube of certain thickness. By pinching the handle, we can control precisely the difference between the infimums of the Log Sobolev functional on two adjacent hand bags. The long round necks serve the following purpose: when two hand bags are joined, the change in the infimum of the Log Sobolev functional happens in a controlled way. In the next few steps we will construct the components inductively in detail.

Step 1. constructing the central component Ω_0. See the figure at the end of the step.

Step 1.1. We start with the standard 3 sphere with three small balls cut out. To be more precise, let

$$D = S^3 - (B_1 \cup B_2 \cup B_3)$$

where S^3 is the standard 3 sphere and $B_i = B(m_i, r)$, $i = 1, 2, 3$, are geodesic balls on S^3 with radius $r > 0$. We take m_1, the center of the ball B_1 at the north pole of S^3; x_2, the center of the ball B_2 at the "left end" of the equator; and x_3, the center of the ball B_3 at the "right end" of the equator. The radius r is so chosen that ∂B_i, $i = 1, 2, 3$, is $h^2 S^2$, the standard 2 sphere with radius h. The radius $h \in (0, 1/4]$ is made sufficiently small so that the following conditions hold:

(1) Lemmas 3.2 and 3.5 hold;

(2) $\lambda(h^2(S^1 \times S^1) \times [-2, 2])) < 0$. That is the infimum of the Log Sobolev functional for the flat tube is negative.

By Lemma 3.6, condition (2) can always be satisfied when h is small enough.

Once chosen, this h will be fixed through out the proof.

Step 1.2. Attach a long round neck $h^2 S^2 \times [0, l]$ to D along ∂B_2 and ∂B_3 respectively. Here $l > 0$ is a large number given by

$$l = \max\{l_0, \frac{1}{a} \ln(1000 A e^{2a}/a^2), \frac{1}{a} \ln(1000 e^{2a} A), 2\}. \tag{3.14}$$

Here l_0, a, A are the numbers in Lemmas 3.2, 3.3 and 3.5. By taking this value for l, all these three lemmas hold and

$$20 A e^{2a} e^{-a(l+k)} \le \frac{1}{2(1+k^2)}, \quad k = 0, 1, 2, 3, ... \tag{3.15}$$

This inequality, to be used shortly in the end of the proof, can be verified easily by finding the maximum of $(1 + k^2) e^{-ak}$.

Let $h^2(S^1 \times S^1) \times S^1 = h^2(S^1 \times S^1) \times [-\pi, \pi]$ be a flat 3 torus, which is regarded as a flat tube given in Definition 3.1. Consider

$$E = h^2(S^1 \times S^1) \times [-\pi, \pi] - B_4.$$

Here $B_4 = B(m_4, h)$ is the geodesic ball of radius h centered at m_4 whose coordinate is $(0, 0, \pi)$. i.e. m_4 is at the bottom of the flat tube. Note h is less than the injectivity radius of the flat torus, which is πh. Therefore we know B_4 is isometric to the Euclidean ball of radius h. Hence $\partial B_4 = h^2 S^2$.

Now we join D with E by a short round neck $h^2 S^2 \times [0, 1]$ by pasting $h^2 S^2 \times \{0\}$ with ∂B_1, and pasting $h^2 S^2 \times \{0\}$ with ∂B_4.

Step 1.3. The metric near the pasted boundaries are smoothed out to satisfy the following conditions.

(1) only the original metric on D near a small neighborhood of ∂B_i, $i = 1, 2, 3$, are perturbed, so that the metric on the attached long round necks stay the same.

(2) only the metric in a small neighborhood of ∂B_4 is perturbed so that the metric on $h^2(S^1 \times S^1) \times [-2, 2]$, which is the top portion of the flat tube, stays intact.

Note the smoothing process is a standard procedure in geometry when one constructs connected sums of two manifolds.

The resulting manifold with boundary is called Ω_0 with metric g_0. By condition (2) in Step 1.1, we have

$$\lambda(\Omega_0, g_0) \le \lambda(h^2(S^1 \times S^1) \times [-2, 2])) < 0. \tag{3.16}$$

For clarity, we write

$$\Omega_0 = Z_0 \cup X \cup H \cup Y_0. \tag{3.17}$$

Here Z_0 is the round neck at the left, which is $h^2 S^2 \times [0, l]$; Y_0 is the round neck at the right, which is $h^2 S^2 \times [0, l]$ again. In order to distinguish the two, we use z to denote points in Z_0, and use y to denote points in Y_0. H denotes the top portion of the flat tube where the third variable of the coordinates is in the interval $[-2, 2]$. i.e. $h^2(S^1 \times S^1) \times [-2, 2]$. We will use the following global coordinate to denote the topological H in the rest of the proof.

$$H = [-\pi, \pi]^2 \times [-2, 2]. \tag{3.18}$$

The metric g_0 on H is just $h^2 g_{S^1 \times S^1} \times g_{R^1}$. The region X is defined to be

$$X = \Omega_0 - (Z_0 \cup H \cup Y_0)$$

We call X the core of Ω_0. The manifold (X, g_0) will serve as the core for all the rest of the components Ω_k.

The shape of Ω_0 is illustrated here.



Step 2. constructing the next component Ω_1 with metric g_1 such that

$$\lambda(\Omega_1, g_1) = \lambda(\Omega_0, g_0) - 1. \tag{3.19}$$

Step 2.1. Attach the round neck $h^2 S^2 \times [0, 1]$ to the left end of Ω_0, forming the round neck $h^2 S^2 \times [0, l+1]$ on the left side, which we call Z_1. Then attach the round neck $h^2 S^2 \times [0, 1]$ to the right end of Ω_0, forming the round neck $h^2 S^2 \times [0, l+1]$ on the right side, which we call Y_1. The resulting domain is called Ω_1 with inherited metric called $\tilde{g}_1$. For convenience we write

$$\Omega_1 = Z_1 \cup X \cup H \cup Y_1.$$

Note $\tilde{g}_1$ is already a smooth metric. In fact $\tilde{g}_1$ is the same as g_0 on X and H, and it is just the product metric on $h^2 g_{S^2} \times g_{R^1}$ on Z_1 and Y_1. But it is not the desired one yet.

Step 2.2. Modify $\tilde{g}_1$ to a new metric g_1 so that (3.19) holds. This modification only happens on H, the top portion of the flat tube. More precisely, this is done by pinching the top portion of the flat tube. Here are the details.

Recall that the top portion of Ω_1 is the flat tube $H = [-\pi, \pi]^2 \times [-2, 2]$. Let θ be a smooth function on Ω_1, satisfying

$$\theta = \theta(x) = \begin{cases} 1, & x \in \Omega_1 - H \\ \text{a number in} \quad (1/2, 1), & x \in H, \quad x \in [-\pi, \pi]^2 \times [-2, -1] \\ 1/2, & x \in H, \quad x \in [-\pi, \pi]^2 \times [-1, 1] \\ \text{a number in} \quad (1/2, 1), & x \in H, \quad x \in [-\pi, \pi]^2 \times [1, 2]. \end{cases}$$

See the figure below.



Now consider the metrics on Ω_1:

$$g_1^{(p)}(x) = \begin{cases} \tilde{g}_1(x), & x \in \Omega_1 - H \\ [\theta^p(x) h^2 g_{S^1 \times S^1}] \times g_{R^1}, & x \in H. \end{cases}$$

We claim that there exists a number $p_1 > 0$ so that

$$\lambda(\Omega_1, g_1^{(p_1)}) = \lambda(\Omega_0, g_0) - 1. \tag{3.20}$$

Here is the proof. Regarding (Ω_0, g_0) as a domain in $(\Omega_1, \tilde{g}_1)$ and applying Lemma 3.5 twice, we know that

$$\lambda(\Omega_1, \tilde{g}_1) \geq \lambda(\Omega_0, g_0) - 2Ae^{-al}$$

for constants $a, A > 0$. By (3.15) with $k = 0$, this leads to

$$\lambda(\Omega_1, \tilde{g}_1) \geq \lambda(\Omega_0, g_0) - 1.$$

Taking $p > 0$ as a variable, the metrics $g_1^{(p)}$ evolves smoothly with p. Lemma 2.5 shows that $\lambda(\Omega_1, g_1^{(p)})$ is a continuous function of p. Observe that

$$\lambda(\Omega_1, g_1^{(0)}) = \lambda(\Omega_1, \tilde{g}_1) \geq \lambda(\Omega_1, g_1) - 1$$

since $g_1^{(0)} = \tilde{g}_1$. By the construction of $g_1^{(p)}$, for $x \in H$ such that $x_3 \in [-1, 1]$,

$$g_1^{(p)}(x) = \left(\frac{1}{2^p} h^2 g_{S^1 \times S^1}\right) \times g_{R^1}.$$

By Lemma 3.6, we know that

$$\lambda(\Omega_1, g_1^{(p)}) \le \lambda(\frac{1}{2^p} h^2 (S^1 \times S^1) \times [-1, 1]) \to -\infty, \quad p \to \infty.$$

By mean value theorem, there exists a number $p = p_1 > 0$ so that (3.20) holds, proving the claim. This metric $g_1^{(p_1)}$ is the desired metric g_1 for Ω_1, satisfying (3.19). This completes the construction of the component (Ω_1, g_1), whose composition is being summarized here for clarity.

$$\Omega_1 = Z_1 \cup X \cup H \cup Y_1. \tag{3.21}$$

where

$$g_1 = \begin{cases} \text{the round metric} \quad h^2 g_{S^2} \times g_{R^1}, & \text{on} \quad Z_1 \cup Y_1 \\ g_0, & \text{on} \quad X \\ \left(\theta^{p_1} h^2 g_{S^1 \times S^1}\right) \times g_{R^1}, & \text{on} \quad H. \end{cases}$$

The shape of Ω_1 is depicted here.



Proceeding inductively, suppose we have constructed

$$\Omega_k = Z_k \cup X \cup H \cup Y_k. \tag{3.22}$$

where

$$g_k = \begin{cases} \text{the round metric} \quad h^2 g_{S^2} \times g_{R^1}, & \text{on} \quad Z_k \cup Y_k \\ g_0, & \text{on} \quad X \\ \left(\theta^{p_k} h^2 g_{S^1 \times S^1}\right) \times g_{R^1}, & \text{on} \quad H, \end{cases}$$

and Z_k and Y_k are round necks of length $l + k$. Now we move to

Step 3. constructing the component Ω_{k+1} so that

$$\lambda(\Omega_{k+1}, g_{k+1}) = \lambda(\Omega_k, g_k) - \frac{1}{k^2 + 1}. \tag{3.23}$$

This is similar to Step 2, with some modification of parameters.

Step 3.1. Attach the round neck $h^2S^2 \times [0,1]$ to the left end of Ω_k, forming the round neck $h^2S^2 \times [0, l+k+1]$ on the left side, which we call Z_{k+1}. Then attach the round neck $h^2S^2 \times [0,1]$ to the right end of Ω_k, forming the round neck $h^2S^2 \times [0, l+k+1]$ on the right side, which we call Y_{k+1}. The resulting domain is called Ω_{k+1} with inherited metric called $\tilde{g}_{k+1}$. i.e.

$$\Omega_{k+1} = Z_{k+1} \cup X \cup H \cup Y_{k+1}.$$

and

$$\tilde{g}_{k+1} = \begin{cases} \text{the round metric} \quad h^2 g_{S^2} \times g_{R^1}, & \text{on} \quad Z_{k+1} \cup Y_{k+1} \\ g_0, & \text{on} \quad X \\ \left(\theta^{p_k} h^2 g_{S^1 \times S^1}\right) \times g_{R^1}, & \text{on} \quad H. \end{cases}$$

Step 3.2. Modify $\tilde{g}_{k+1}$ to a new metric g_{k+1} so that (3.23) holds.
This is again done by pinching H, the top portion of the flat tube. Here are the details. Let θ be the smooth function as in Step 2. Now consider the metrics on Ω_{k+1}:

$$g^{(p)}_{k+1}(x) = \begin{cases} \tilde{g}_{k+1}(x), & x \in \Omega_{k+1} - H \\ \left(\theta^p(x) h^2 (S^1 \times S^1)\right) \times g_{R^1}, & x \in H. \end{cases}$$

We claim that there exists a number $p_{k+1} > 0$ so that

$$\lambda(\Omega_{k+1}, g^{(p_{k+1})}_{k+1}) = \lambda(\Omega_k, g_k) - \frac{1}{k^2+1}. \tag{3.24}$$

Here is the proof. Regarding (Ω_k, g_k) as a domain in $(\Omega_{k+1}, \tilde{g}_{k+1})$ and applying Lemma 3.5 twice, we know that

$$\lambda(\Omega_{k+1}, \tilde{g}_{k+1}) \geq \lambda(\Omega_k, g_k) - 2Ae^{-a(l+k)}$$

for constants $a, A > 0$. Note the length of Z_k and Y_k are $k+l$, which explains the appearance of the exponential term $e^{-a(l+k)}$. By (3.15), this leads to

$$\lambda(\Omega_{k+1}, \tilde{g}_{k+1}) \geq \lambda(\Omega_k, g_k) - \frac{1}{k^2+1}.$$

Taking $p > 0$ as a variable, the metrics $g^{(p)}_{k+1}$ evolves smoothly with p. Lemma 2.5 shows that $\lambda(\Omega_{k+1}, g^{(p)}_{k+1})$ is a continuous function of p. Observe that

$$\lambda(\Omega_{k+1}, g^{(p_k)}_{k+1}) = \lambda(\Omega_{k+1}, \tilde{g}_{k+1}) \geq \lambda(\Omega_k, g_k) - \frac{1}{k^2+1}$$

since $g^{(p_k)}_{k+1} = \tilde{g}_{k+1}$. By the construction of $g^{(p)}_{k+1}$, for $x \in H$ such that $x_3 \in [-1,1]$,

$$g^{(p)}_{k+1}(x) = \frac{1}{2^p} h^2 g_{S^1 \times S^1} \times g_{R^1}.$$

From Lemma 3.6, we know that

$$\lambda(\Omega_{k+1}, g^{(p)}_{k+1}) \leq \lambda(\frac{1}{2^p} h^2 (S^1 \times S^1) \times [-1,1]) \to -\infty, \quad p \to \infty.$$

By mean value theorem, there exists a number $p = p_{k+1} \geq p_k$ so that (3.24) holds, proving the claim. This metric $g^{(p_{k+1})}_{k+1}$ is the desired metric g_{k+1} for Ω_{k+1}, satisfying (3.23). This completes the construction of the component (Ω_{k+1}, g_{k+1}), finishing the induction. To summarize,

$$\Omega_{k+1} = Z_{k+1} \cup X \cup H \cup Y_{k+1}. \tag{3.25}$$

and

$$g_{k+1} = \begin{cases} \text{the round metric} \quad h^2 g_{S^2} \times g_{R^1}, & \text{on} \quad Z_{k+1} \cup Y_{k+1} \\ g_0, & \text{on} \quad X \\ \left(\theta^{p_{k+1}} h^2 g_{S^1 \times S^1}\right) \times g_{R^1}, & \text{on} \quad H. \end{cases}$$

The shape of Ω_{k+1} is depicted here.



Step 4. pasting together the components to form the manifold $\mathbf{M}$. See the figure at the end of the step.

In the last step, we have constructed the manifolds (Ω_k, g_k) for $k = 0, 1, 2, 3, ...$ Now we define,

$$(\Omega_{-k}, g_{-k}) = (\Omega_k, g_k), \quad k = 1, 2, ...$$

Finally, we take

$$\mathbf{M} = \cup_{k=-\infty}^{\infty} \Omega_k \tag{3.26}$$

which is the connected, non-overlapping union of Ω_k, for all integers k in the following pattern. We connect Ω_k with Ω_{k+1} by pasting the right end of Y_k with left end of Z_{k+1}. Here $k = ..., -2, -1, 0, 1, 2, ...$ The metric on $\mathbf{M}$, which is inherited from g_k, is denoted by g.

It is clear that $\mathbf{M}$ is a complete, connected manifold. Now let us prove $\mathbf{M}$ has bounded geometry. Note that except for the top portions of Ω_k, which is denoted by H, the manifold $\mathbf{M}$ is consisted of round necks or flat tubes of fixed aperture. Hence we just need to prove that (H, g) has bounded geometry. The metric g on $H \subset \Omega_k$ is given by $(\theta^{p_k}(x) h^2 g_{S^1 \times S^1}) \times g_{R^1}$, where $\theta(x) = 1/2$ when $x_3 \in [-1, 1]$ and $1/2 \le \theta \le 1$. Write

$$\lambda_k = \lambda(\Omega_k, g), \quad k = 0, 1, 2, ...$$

Recall by construction that

$$\lambda_0 - \Sigma_{j=1}^{|k|} \frac{1}{j^2} = \lambda_k \le 0, \quad |k| = 1, 2, ...,$$

which implies

$$\lambda\left((1/2)^{p_k} h^2 (S^1 \times S^1) \times [-1, 1]\right) \ge \lambda_k \ge \lambda_0 - 10.$$

If $\{p_k\}$ is unbounded, then by Lemma 3.6, the left hand side of the above inequality tends to $-\infty$ when $k \to \infty$, which leads to a contradiction. Hence $\{p_k\}$ is a bounded sequence of positive numbers. Since θ is a smooth bounded function, we know θ^{p_k} has uniformly bounded C^∞ norm. Therefore we have proven that $\mathbf{M}$ has bounded geometry everywhere. The shape of $\mathbf{M}$ is depicted here.



Step 5. proving that the Log Sobolev functional on $\mathbf{M}$ does not have an extremal.

We use the method of contradiction. Suppose that a smooth function v, $\|v\|_{L^2(\mathbf{M})} = 1$, is an extremal for the Log Sobolev functional whose infimum is $\lambda = \lambda(\mathbf{M}, g)$. Then

$$\lambda = \int_{\mathbf{M}} (4|\nabla v|^2 + Rv^2 - v^2 \ln v^2) dg$$

and v is a smooth solution to equation (2.1) i.e.

$$4\Delta v - Rv + 2v \ln v + \lambda v = 0.$$

Let us recall that

$$\Omega_k = Z_k \cup X \cup H \cup Y_k,$$

where $Z_k = h^2 S^2 \times [0, l+k]$ and $Y_k = h^2 S^2 \times [0, l+k]$ are round necks on the left and right side of the core X respectively. In order to distinguish these two necks, we use z to denote points in Z_k with a coordinate $z = (z_1, z_2, z_3)$ described in Definition 3.1; and likewise we use y to denote points in Y_k with a coordinate $y = (y_1, y_2, y_3)$ described in Definition 3.1. These two coordinates are regarded as independent ones.

For each $k = 1, 2, \ldots$, we construct a cut-off function $\eta_k \in W_0^{1,\infty}(\Omega_k)$ as follows.

$$\eta_k = \begin{cases} \eta_k(z) = z_3, & z \in Z_k, \quad 0 \le z_3 \le 1, \\ \eta_k(z) = 1, & z \in Z_k, \quad 1 \le z_3 \le l+k, \\ \eta_k(x) = 1, & x \in X \cup H, \\ \eta_k(y) = 1, & y \in Y_k, \quad 0 \le y_3 \le l+k-1, \\ \eta_k(y) = 1 - (y_3 - l - k + 1), & y \in Y_k, \quad l+k-1 \le y_3 \le l+k. \end{cases} \tag{3.27}$$

The following figure depicts the definition of η_k.



Since v solves (2.1), we can apply Lemma 3.4 by taking $E = \Omega_k$ and $F = \mathbf{M}$ there to get

$$\lambda_k \int (v\eta_k)^2 dg \le \lambda \int (v\eta_k)^2 dg + 4 \int v^2 |\nabla \eta_k|^2 dg - \int (v\eta_k)^2 \ln \eta_k^2 dg.$$

Here $\lambda_k = \lambda(\Omega_k, g_k)$. Observe that $|\nabla \eta_k| \le 1$ and that the function $(\eta_k)^2 \ln \eta_k^2$, which is nonzero only in the support of $\nabla \eta_k$, is bounded from below by $-e^{-1}$. Therefore

$$\lambda_k \int (v\eta_k)^2 dg \le \lambda \int (v\eta_k)^2 dg + 5 \int_{supp \nabla \eta_k} v^2 dg. \tag{3.28}$$

By definition of η_k, $supp \nabla \eta_k$ is the disjoint union of two short round necks, i.e.

$$supp \nabla \eta_k = Z_{k1} \cup Y_{k1} \tag{3.29}$$

where

$$Z_{k1} \equiv \{z \in Z_k \, | 0 \le z_3 \le 1\}, \qquad Y_{k1} \equiv \{y \in Y_k \, | l + k - 1 \le y_3 \le l + k\}.$$

Hence (3.28) implies

$$(\lambda_k - \lambda) \int (v\eta_k)^2 dg \le 5 \int_{Z_{k1}} v^2 dg + 5 \int_{Y_{k1}} v^2 dg = 5 \int_{supp \nabla \eta_k} v^2 dg. \tag{3.30}$$

Next we prove that the right hand side of (3.30) is exponentially small. Observe that Z_{k1} is a middle segment of $Y_{k-1} \cup Z_k$, which is, when writing in one coordinate, a round neck of the form $h^2 S^2 \times [0, 2l + 2k - 1]$. The segments

$$W_{k-1} \equiv \{y \in Y_{k-1} | 0 \le y_3 \le 2\} \quad \text{and} \quad E_k \equiv \{z \in Z_k | k + l - 2 \le z_3 \le k + l\}$$

are at the left and right end of the round neck respectively. See the figure below.



By Lemma 3.2, we have

$$\int_{Z_{k1}} v^2 dg \le A e^{-a(l+k-1)} \left[\int_{\{y \in Y_{k-1} | 0 \le y_3 \le 2\}} v^2 dg + \int_{\{z \in Z_k | k+l-2 \le z_3 \le k+l\}} v^2 dg \right].$$

Note from (3.27) that

$$\eta_{k-1} = 1 \quad \text{in} \quad W_{k-1} = \{y \in Y_{k-1} | 0 \le y_3 \le 2\} \subset Y_{k-1} \subset \Omega_{k-1},$$

$$\eta_k = 1 \quad \text{in} \quad E_k = \{z \in Z_k | k+l-1 \le z_3 \le k+l\} \subset Z_k \subset \Omega_k.$$

Hence

$$\int_{Z_{k1}} v^2 dg \le A e^{-a(l+k-1)} \left[\int_{\Omega_{k-1}} (\eta_{k-1} v)^2 dg + \int_{\Omega_k} (\eta_k v)^2 dg \right]. \tag{3.31}$$

Similarly, we see that Y_{k1} is a middle segment of $Y_k \cup Z_{k+1}$, which is, when writing in one coordinate, a round neck of the form $h^2 S^2 \times [0, 2l + 2k + 1]$. The segments

$$W_k \equiv \{y \in Y_k | 0 \le y_3 \le 2\} \quad \text{and} \quad E_{k+1} \equiv \{z \in Z_{k+1} | k+l-1 \le z_3 \le k+l+1\}$$

are the left and right end of the round neck. See the figure below.



By Lemma 3.2, we have

$$\int_{Y_{k1}} v^2 dg \le A e^{-a(l+k)} \left[\int_{\{y \in Y_k | 0 \le y_3 \le 2\}} v^2 dg + \int_{\{z \in Z_{k+1} | k+l-1 \le z_3 \le k+l+1\}} v^2 dg \right].$$

Note that

$$\eta_k = 1 \quad \text{in} \quad W_k = \{y \in Y_k | 0 \le y_3 \le 2\} \subset Y_k \subset \Omega_k$$

and

$$\eta_{k+1} = 1 \quad \text{in} \quad E_{k+1} = \{z \in Z_{k+1} | k+l-1 \le z_3 \le k+l+1\} \subset Z_{k+1} \subset \Omega_{k+1}.$$

Hence

$$\int_{Y_{k1}} v^2 dg \le A e^{-a(l+k)} \left[\int_{\Omega_k} (\eta_k v)^2 dg + \int_{\Omega_{k+1}} (\eta_{k+1} v)^2 dg \right]. \tag{3.32}$$

By this, (3.31) and (3.29), we obtain

$$\int_{supp \nabla \eta_k} v^2 dg \le A e^{-a(l+k-1)} \left[\int_{\Omega_{k-1}} (\eta_{k-1} v)^2 dg + 2 \int_{\Omega_k} (\eta_k v)^2 dg + \int_{\Omega_{k+1}} (\eta_{k+1} v)^2 dg \right]$$

where $k = 1, 2, 3, ...$ Recall that $(\Omega_{-k}, g_{-k}) = (\Omega_k, g_k)$ be definition. Therefore, we can derive, in a similar manner,

$$\int_{supp\nabla\eta_k} v^2 dg \le A e^{-a(l+|k|-1)} \left[\int_{\Omega_{k-1}} (\eta_{k-1} v)^2 dg + 2 \int_{\Omega_k} (\eta_k v)^2 dg + \int_{\Omega_{k+1}} (\eta_{k+1} v)^2 dg \right]$$

where $k = 0, -1, -2, -3, ...$ Adding the last two inequalities together, we deduce

$$\begin{aligned} \Sigma_{k=-\infty}^{\infty} \int_{supp\nabla\eta_k} v^2 dg &= \Sigma_{k=-\infty}^{\infty} \int_{Z_{k1}} v^2 dg + \Sigma_{k=-\infty}^{\infty} \int_{Y_{k1}} v^2 dg \\ &\le 4Ae^{2a} \Sigma_{k=-\infty}^{\infty} e^{-a(l+|k|)} \int_{\Omega_k} (v\eta_k)^2 dg. \end{aligned}$$

By (3.30), this implies

$$\Sigma_{k=-\infty}^{\infty} (\lambda_k - \lambda - 20Ae^{2a} e^{-a(l+k)}) \int (v\eta_k)^2 dg \le 0.$$

Recall, by construction,

$$\lambda_k - \lambda \ge \lambda_k - \lambda_{k+1} = \frac{1}{k^2+1}, \quad k = 1, 2, 3, ...$$

and

$$\lambda_k - \lambda \ge \lambda_k - \lambda_{k-1} = \frac{1}{k^2+1}, \quad k = -1, -2, -3, ...,$$

and

$$\lambda_0 - \lambda \ge \lambda_0 - \lambda_1 = 1.$$

So finally we deduce

$$\Sigma_{k=-\infty}^{\infty} (\frac{1}{1+k^2} - 20Ae^{2a} e^{-a(l+|k|)}) \int (v\eta_k)^2 dg \le 0.$$

This is a contradiction because $\frac{1}{1+k^2} - 20Ae^{2a}e^{-a(l+|k|)} > 0$ by our choice of l in (3.15). Therefore no extremal for the Log Sobolev functional exists. □

4. *W* ENTROPY AND A NO BREATHER RESULT FOR NONCOMPACT RICCI FLOW

In this section we discuss some applications of Theorem 1.1 to Perelman's W entropy and Hamilton's Ricci flow. We will use the following notations. $g = g(t)$ is a metric which evolves with time; $d(x, y, t)$ or $d(x, y, g(t))$ will denote the corresponding distance function; $dg(t)$ denotes the volume element under $g(t)$; We will still use ∇, Δ the corresponding gradient and Laplace-Beltrami operator, when no confusion arises.

The following definition is one of several equivalent ways in which Perelman's W entropy can be written.

Definition 4.1. *(W entropy) Let* $v \in W^{1,2}(\mathbf{M})$ *and* $\tau > 0$ *be a parameter. The* W *entropy is the quantity*

$$W(g, v, \tau) \equiv \int_{\mathbf{M}} \left[\tau(4|\nabla v|^2 + Rv^2) - v^2 \ln v^2 - \frac{n}{2}(\ln 4\pi\tau)\ v^2 - nv^2 \right] dg. \tag{4.1}$$

Let $c > 0$ be a positive constant, it is clear that the W entropy has the following scaling invariant property

$$W(cg, c^{-n/2}v, c\tau) = W(g, v, \tau).$$

Hence we can always take $\tau = 1$ if necessary. If $\tau = 1$ and $\|v\|_{L^2(\mathbf{M})} = 1$, then

$$\begin{aligned} W(g, v, 1) &= \int_{\mathbf{M}} \left[(4|\nabla v|^2 + Rv^2) - v^2 \ln v^2\right] dg - \frac{n}{2}(\ln 4\pi) - n \\ &= L(v, g) - \frac{n}{2}(\ln 4\pi) - n. \end{aligned} \tag{4.2}$$

Here $L(v, g)$ is the Log Sobolev functional given in (1.1). Therefore, the W entropy and the Log Sobolev functional differ only by a normalizing constant after scaling.

Perelman also introduced the so called μ invariant.

Definition 4.2. *Given a noncompact manifold* $(\mathbf{M}, g)$ *and parameter* $\tau > 0$*, the* μ *invariant is the quantity*

$$\mu(g, \tau) = \inf\{W(g, v, \tau) \,|\, v \in C_0^\infty(\mathbf{M}),\ \|v\|_{L^2(\mathbf{M})=1}\}.$$

In view of Definition 1.1, we introduce μ invariant near infinity.

Definition 4.3. *Given a noncompact manifold* $(\mathbf{M}, g)$ *and parameter* $\tau > 0$*, the* μ *invariant at infinity is the quantity*

$$\begin{aligned} \mu_\infty(g, \tau) = \lim_{r\to\infty} \inf\{\int_{\mathbf{M}-B(0,r)} &\left[\tau(4|\nabla v|^2 + Rv^2) - v^2 \ln v^2 - \frac{n}{2}(\ln 4\pi\tau)\ v^2 - nv^2\right] dg \,| \\ &v \in C_0^\infty(\mathbf{M} - B(0, r)), \quad \|v\|_{L^2(\mathbf{M}-B(0,r))=1}\}. \end{aligned}$$

Since the W entropy and the Log Sobolev functional differ only by a constant after scaling, Theorem 1.1 can be immediately transplanted as

Theorem 4.1. *(a). Let* $\mathbf{M}$ *be a complete, connected noncompact manifold with bounded geometry, and* $\tau > 0$ *be a parameter. Suppose* $\mu(\mathbf{M}, \tau) < \lambda_\infty(\mathbf{M}, \tau)$*, then there exists a smooth extremal* v *for the* W *entropy (4.1). Also, there exist positive constants* $a, A > 0$ *and a point* $0 \in \mathbf{M}$ *such that*

$$v(x) \le Ae^{-ad^2(x,0)}.$$

(b). There exists a complete, connected noncompact manifold with bounded geometry such that $\mu(\mathbf{M}, \tau) < \lambda_\infty(\mathbf{M}, \tau)$*, but the* W *entropy (4.1) does not have an extremal.*

In the rest of the section, we describe two more applications of this theorem. The first one is an extension of Perelman's monotonicity formula for the W entropy from the compact case to some noncompact ones.

Let us briefly recall Perelman's monotonicity formula. Consider the final value problem of the conjugate heat equation coupled with the Ricci flow $(\mathbf{M}, g(t))$ on a compact manifold $\mathbf{M}$ and on the time interval $[t_1, t_2]$.

$$\begin{cases} \Delta u - Ru + u_t = 0, \quad t \in [t_1, t_2] \\ u(x, t_2) = u_2 \\ \partial_t g(t) = -2Ric, \quad t \in [t_1, t_2]. \end{cases} \tag{4.3}$$

Here Δ is the Laplace-Beltrami operator with respect to the metric $g(t)$; R and Ric are the scalar curvature and Ricci curvature with respect to $g(t)$; and $u_2 = u_2(x)$ is a smooth function such that $\|u_2\|_{L^1(\mathbf{M},g(t_2))} = 1$. In the definition of the W entropy, we take $\tau = L - t$ and $v(\cdot, t) = \sqrt{u(\cdot, t)}$. Perelman ([P] section 3) proved that

$$\frac{d}{dt} W(g(t), v(\cdot, t), L - t) = 2\tau \int_{\mathbf{M}} \left| Ric - Hess \ln u - \frac{1}{2\tau} g \right|^2 u \, dg(t). \tag{4.4}$$

If $\mathbf{M}$ is noncompact, then the above formula needs certain justification. One reason is that the term $Hess \ln u$ may grow to infinity and hence the integral may diverge. Consequently, certain extra decay conditions are needed on u and $Hess \ln u$. When u is the fundamental solution of the conjugate heat equation, a noncompact version of the above formula has been carefully established in [CCGGIIKLLN3] Chapters 19, 20, 21 and the paper [CTY]. They employed a number of technical tools such as Log gradient bounds for positive solutions of (4.3) and pointwise bounds on the fundamental solution of (4.3). With the help of these tools and the decay estimate of extremals of the W entropy, we extend (4.4) to a noncompact case where the final value u_2 is the square of an extremal of the W entropy. The point of the following corollary is that once an extremal exists, then no other decay conditions are needed.

Corollary 4.1. *Let $(\mathbf{M}, g(t))$ be a Ricci flow which has bounded geometry in the finite time interval $[t_1, t_2]$. Assume also that the 4-th order derivatives of the curvature tensor are uniformly bounded in $\mathbf{M} \times [t_1, t_2]$. Let $\tau = L - t$ with $L > t_2$ be a parameter. Suppose the W entropy $W(g(t_2), v, T - t_2)$ has an extremal v_2. Let u be the solution of the final value problem of the conjugate heat equation:*

$$\begin{cases} \Delta u - Ru + u_t = 0, & t \in [t_1, t_2] \\ u(x, t_2) = v_2^2 \\ \partial_t g(t) = -2Ric, & t \in [t_1, t_2]. \end{cases}$$

Let $v = v(x, t) = \sqrt{u(x, t)}$. Then, for all $t \in [t_1, t_2]$, the W entropy $W(g(t), v, T - t)$ is well defined. Moreover

$$\frac{d}{dt} W(g(t), v, L - t) = 2\tau \int_{\mathbf{M}} \left| Ric - Hess \ln u - \frac{1}{2\tau} g \right|^2 u \, dg(t).$$

Proof.

The task is to show that relevant integrands has quadratic exponential decay at infinity. After this, the proof is the same as Perelman's in the compact case.

Step 1. First we show that there exist positive constants A_1, a_1 and a point $0 \in \mathbf{M}$ such that

$$u(x, t) \le A_1 e^{-a_1 d^2(x, 0, t)}. \tag{4.5}$$

This bound follows from the decay of the extremal v_2 in Theorem 4.1 (a) and the following bounds on $G = G(x, t; y, t_2)$, the fundamental solution of the conjugate heat equation (4.3). Observe that the Ricci flow has bounded geometry in the finite time interval $[t_1, t_2]$. Hence the distance functions $d(x, 0, t)$ are equivalent when $t \in [t_1, t_2]$. The same can be said for volumes $|B(x, r, t)|_{g(t)}$. By [CCGGIIKLLN3] Chapters 19 or [CTY] Section 5, there are the bounds:

$$G(x,t;y,t_2) \ge \frac{1}{\alpha\sqrt{|B(x,\sqrt{t_2-t},t)|_{g(t)}}\sqrt{|B(y,\sqrt{t_2-t},t)|_{g(t)}}} e^{-\frac{d^2(x,y,t)}{\beta(t_2-t)}},$$

$$G(x,t;y,t_2) \le \frac{\alpha}{\sqrt{|B(x,\sqrt{t_2-t},t)|_{g(t)}}\sqrt{|B(y,\sqrt{t_2-t},t)|_{g(t)}}} e^{-\beta\frac{d^2(x,y,t)}{(t_2-t)}},$$

where the constants α and β depend on $\mathbf{M}$, t_1 and t_2. These bounds can be regarded as generalization of the bounds in [LY] for the heat equation under fixed metrics. By the assumption of bounded geometry and classical volume comparison theorem, there exist positive constants c, c_1 and c_2 such that

$$c_1\min\{1,(t_2-t)^{n/2}\} \le |B(x,\sqrt{t_2-t},t)|_{g(t)} \le c(t_2-t)^{n/2}e^{c_2\sqrt{t_2-t}},$$

$$c_1\min\{1,(t_2-t)^{n/2}\} \le |B(y,\sqrt{t_2-t},t)|_{g(t)} \le c(t_2-t)^{n/2}e^{c_2\sqrt{t_2-t}},$$

Hence we have the bounds: for $t\in[t_1,t_2]$ and $x,y\in\mathbf{M}$,

$$\frac{1}{\alpha(t_2-t)^{n/2}}e^{-\beta\frac{d^2(x,y,t)}{(t_2-t)}} \le G(x,t;y,t_2) \le \frac{\alpha}{(t_2-t)^{n/2}}e^{-\beta\frac{d^2(x,y,t)}{(t_2-t)}}, \tag{4.6}$$

where the constant $\alpha=\alpha(\mathbf{M},t_1,t_2)$ may have changed from its previous value. Therefore,

$$u(x,t)=\int_{\mathbf{M}}G(x,t;y,t_2)u(y,t_2)dg(t_2) \le \int_{\mathbf{M}}\frac{\alpha}{(t_2-t)^{n/2}}e^{-\beta\frac{d^2(x,y,t)}{(t_2-t)}}u(y,t_2)dg(t_2)$$

By Theorem 4.1 (a) (in fact Lemma 2.3 is sufficient), there exist positive constants $a, A>0$ such that

$$u(x,t_2)=v_2^2(x)\le 2Ae^{-2ad^2(x,0,t_2)}. \tag{4.7}$$

The last two inequalities imply

$$u(x,t)\le 2\alpha A\int_{\mathbf{M}}\frac{1}{(t_2-t)^{n/2}}e^{-\beta\frac{d^2(x,y,t)}{(t_2-t)}}e^{-2ad^2(y,0,t_2)}dg(t_2). \tag{4.8}$$

By triangle inequality, there exist $a_1>0$ such that

$$-\beta\frac{d^2(x,y,t)}{(t_2-t)}-2ad^2(y,0,t_2)\le -a_1d^2(x,0,t_2)-\beta\frac{d^2(x,y,t)}{2(t_2-t)}.$$

Here we used the fact that distances at different time levels are comparable again. Hence

$$\begin{aligned}
&\int_{\mathbf{M}}\frac{1}{(t_2-t)^{n/2}}e^{-\beta\frac{d^2(x,y,t)}{(t_2-t)}}e^{-2ad^2(y,0,t_2)}dg(t_2)\\
&\le e^{-a_1d^2(x,0,t_2)}\int_{\mathbf{M}}\frac{1}{(t_2-t)^{n/2}}e^{-\beta\frac{d^2(x,y,t)}{2(t_2-t)}}dg(t_2)\\
&= e^{-a_1d^2(x,0,t_2)}[\Sigma_{k=0}^{\infty}\int_{2^{k-1}\sqrt{t_2-t}\le d(x,y,t)\le 2^k\sqrt{t_2-t}}\frac{1}{(t_2-t)^{n/2}}e^{-\beta\frac{d^2(x,y,t)}{2(t_2-t)}}dg(t_2)\\
&\qquad +\int_{d(x,y,t)\le\sqrt{t_2-t}}\frac{1}{(t_2-t)^{n/2}}e^{-\beta\frac{d^2(x,y,t)}{2(t_2-t)}}dg(t_2)].
\end{aligned}$$

Since $\mathbf{M}$ has bounded geometry, the classical volume comparison theorem tells us

$$|B(x, 2^k\sqrt{t_2 - t}, t)|_{g(t_2)} \le Ce^{c2^k\sqrt{t_2-t}}(t_2 - t)^{n/2}.$$

Here we just used the fact that volume elements at different time levels in $[t_1, t_2]$ are equivalent. Hence

$$\begin{aligned}&\int_{\mathbf{M}} \frac{1}{(t_2 - t)^{n/2}} e^{-\beta\frac{d^2(x,y,t)}{(t_2-t)}} e^{-2ad^2(y,0,t_2)} dg(t_2)\\ &\le e^{-a_1 d^2(x,0,t_2)} \left[\Sigma_{k=0}^{\infty} Ce^{c2^k\sqrt{t_2-t}} e^{-\beta 2^{2(k-1)}/2} + Ce^{c\sqrt{t_2-t_1}}\right],\end{aligned}$$

which shows

$$\int_{\mathbf{M}} \frac{1}{(t_2 - t)^{n/2}} e^{-\beta\frac{d^2(x,y,t)}{(t_2-t)}} e^{-2ad^2(y,0,t_2)} dg(t_2) \le Ce^{-a_1 d^2(x,0,t_2)}, \tag{4.9}$$

where C depends on $t_2 - t_1$. Substituting this to (4.8), we deduce

$$u(x, t) \le A_1 e^{-a_1 d^2(x,0,t_2)}.$$

This proves the bound in (4.5).

Step 2. We prove that the integrand in the W entropy has quadratic exponential decay.

For convenience, we denote the integrand in the W entropy as

$$i(u) = i(u)(x, t) \equiv \left[\tau(\frac{|\nabla u|^2}{u} + Ru) - u \ln u - \frac{n}{2} \ln(4\pi\tau)u - nu\right](x, t). \tag{4.10}$$

Here we have used the relation that $u = v^2$ on (4.1). We now prove that there exist positive constants A_1 and a_1 such that

$$|i(u)(x, t)| \le A_1 e^{-a_1 d^2(x,0,t)}. \tag{4.11}$$

By the bound (4.5), we know that the term $u \ln u$ satisfies

$$|u \ln u| = \sqrt{u}\sqrt{u}|\ln u| \le C\sqrt{u} \le A_1 e^{-a_1 d^2(x,0,t)},$$

whence it also has quadratic exponential decay. Here the values of A_1 and a_1 may have changed. So it suffices to prove that the term $\frac{|\nabla u|^2}{u}$ has the decay too.

To this end, we recall by direct computation (see Proposition 6.1.2 in [Z] e.g.) that

$$\begin{aligned}&H^*(\frac{|\nabla u|^2}{u} + Ru)\\ &= \frac{2}{u}\left(u_{ij} - \frac{u_i u_j}{u}\right)^2 + 2\nabla R\nabla u + \frac{4}{u} Ric(\nabla u, \nabla u) + 2|Ric|^2 u + 2\nabla R\nabla u + 2u\Delta R.\end{aligned}$$

Here $H^* = \Delta - R + \partial_t$ is the conjugate heat operator. Thus

$$H^*(\frac{|\nabla u|^2}{u} + Ru) \ge -K_1(|\nabla u| + |u| + \frac{|\nabla u|^2}{u}),$$

where the constant $K_1(\ge 0)$ depends on the supremum of $|\nabla R|$, $|\Delta R|$ and the lower bound of Ric. Since $|\nabla u| \le \frac{|\nabla u|^2}{u} + u$, we deduce

$$H^*(\frac{|\nabla u|^2}{u} + Ru) \ge -K_1(\frac{|\nabla u|^2}{u} + Ru) - K_2 u, \tag{4.12}$$

where K_2 depends on K_1, the supremum of $|R|$ and u. We mention that all the curvatures involved here are bounded according to our assumption on the Ricci flow.

At time t_2, $u(x,t_2)=v_2^2$. Hence $\frac{|\nabla u|^2}{u}+Ru=4|\nabla v_2|^2+Rv_2^2$. Notice that v_2 satisfies the equation for extremals: for $\tau=L-t_2$,

$$\tau(4\Delta v_2-Rv_2)+2v_2\ln v_2+\frac{n}{2}\ln(4\pi\tau)v_2+nv_2+\mu v_2=0$$

By Lemma 2.1 part (b), we have

$$\sup_{B(x,1/2,t_2)}|\nabla v_2|^2\le C\int_{B(x,1,t_2)}v_2^2dg(t_2)\le CA^2e^{-2ad^2(x,0,t_2)},$$

where the last inequality is due to the decay of v_2 in (4.7). By this and the decay of v_2 again, we know that, at time t_2,

$$\left|\frac{|\nabla u|^2}{u}+Ru\right|(x,t_2)\le A_1e^{-a_1d^2(x,0,t_2)}. \tag{4.13}$$

Now define

$$Q=Q(u)=e^{K_1t}\left(\frac{|\nabla u|^2}{u}+Ru\right).$$

By (4.12) and (4.13), we know that

$$\begin{cases}\Delta Q-RQ+\partial_tQ\ge -K_2e^{K_1t}u, & t\in[t_1,t_2],\\ Q(\cdot,t_2)\le A_1e^{K_1t_2}e^{-a_1d^2(x,0,t_2)}.\end{cases}$$

By the maximum principle (see [CCGGIIKLLN2] Chapter 12 e.g.), this implies, for $t\in[t_1,t_2]$,

$$\begin{aligned}Q(x,t)\le&\int_{\mathbf{M}}G(x,t;y,t_2)A_1e^{K_1t_2}e^{-a_1d^2(y,0,t_2)}dg(t_2)\\&+\int_t^{t_2}\int_{\mathbf{M}}G(x,t;y,s)K_2e^{K_1t_2}u(y,s)dg(s)ds.\end{aligned}\tag{4.14}$$

We mention that even though Q is a smooth function, it may not be a bounded one for each time level, due to the appearance of the term $\frac{|\nabla u|^2}{u}$. In order to apply the maximum principle, one needs some growth condition on Q near infinity. The way to justify (4.14) is to replace u by the function u_ϵ which is the solution to

$$\begin{cases}\Delta u_\epsilon-Ru_\epsilon+\partial_tu_\epsilon=0, & t\in[t_1,t_2]\\ u_\epsilon(x,t_2)=v_2^2+\epsilon\\ \partial_tg(t)=-2Ric, & t\in[t_1,t_2].\end{cases}$$

Here $\epsilon>0$ is a positive number. It is clear that $u_\epsilon\to u$ pointwise when $\epsilon\to 0$. Since u_ϵ is bounded from above and below by positive constants, we know that $Q_\epsilon=Q(u_\epsilon)$ is a bounded function. Moreover it holds

$$\begin{cases}\Delta Q_\epsilon-RQ_\epsilon+\partial_tQ_\epsilon\ge -K_2e^{K_1t}u_\epsilon, & t\in[t_1,t_2],\\ Q_\epsilon(\cdot,t_2)\le A_1e^{K_1t_2}e^{-a_1d^2(x,0,t_2)}+C\epsilon.\end{cases}$$

Now we can apply the maximum principle for Q_ϵ to derive

$$\begin{aligned} Q_\epsilon(x,t) \le &\int_{\mathbf{M}} G(x,t;y,t_2) Q_\epsilon(y,t_2) dg(t_2) \\ &+ \int_t^{t_2} \int_{\mathbf{M}} G(x,t;y,s) K_2 e^{K_1 t_2} u_\epsilon(y,s) dg(s) ds. \end{aligned} \tag{4.15}$$

Taking $\epsilon \to \infty$, this implies (4.14).

By (4.14), (4.6) and (4.5), we derive

$$\begin{aligned} Q(x,t) \le A_1 e^{K_1 t_2} &\int_{\mathbf{M}} \frac{\alpha}{(t_2-t)^{n/2}} e^{-\beta \frac{d^2(x,y,t)}{(t_2-t)}} e^{-a_1 d^2(y,0,t_2)} dg(t_2) \\ &+ K_2 e^{K_1 t_2} \int_t^{t_2} \int_{\mathbf{M}} \frac{\alpha}{(s-t)^{n/2}} e^{-\beta \frac{d^2(x,y,t)}{(s-t)}} A_1 e^{-a_1 d^2(y,0,s)} dg(s) ds. \end{aligned}$$

Using the fact that distance functions and volume elements at different time levels are equivalent, we can apply (4.9) to the above inequality to deduce

$$Q(x,t) \le C e^{-cd^2(x,0,t)}$$

where c and C are positive constants which may depend on t_1 and t_2. This proves that

$$\left| \frac{|\nabla u|^2}{u} + Ru \right| (x,t) \le C e^{-cd^2(x,0,t)}, \quad t \in [t_1, t_2],$$

which implies (4.11).

Step 3. Completion of the proof.

Let u and τ be the same as in the statement of the Corollary. In the paper [P] Proposition 9.1, Perelman introduced the quantity

$$P(u) = \tau(-2\Delta u + \frac{|\nabla u|^2}{u} + Ru) - u \ln u - \frac{n}{2} \ln(4\pi\tau) u - nu \tag{4.16}$$

and proved that

$$H^* P(u) = 2\tau \left| Ric - Hess \ln u - \frac{g}{2\tau} \right|^2 u. \tag{4.17}$$

We mention that in [P], the quantity $P(u)$ here is denoted by $v = v(f)$ where f is determined by $u = \frac{e^{-f}}{(4\pi\tau)^{n/2}}$. Observe that

$$P(u) = -2\tau \Delta u + i(u) \tag{4.18}$$

where $i(u)$ is the integrand of the W entropy used in the previous step.

Next we will integrate (4.17). However, at the moment, we do not know the if terms involved are integrable. So we need to use certain cut off function. Let $L = L(x)$ be a smooth function on $\mathbf{M}$ such that

$$\begin{aligned} &|\nabla L(x)| + |\nabla^2 L(x)| + |\nabla^3 L(x)| + |\nabla^4 L(x)| \le C_1, \qquad x \in \mathbf{M}, \\ &C_1^{-1} L(x) \le d(x, 0, g(t_1)) \le C_1 L(x), \qquad x \in \mathbf{M}. \end{aligned}$$

Here the covariant derivatives are with respect to $g(t_1)$. Under our assumption of bounded geometry, it is well known that such a function exists. See for example Proposition 19.37 and the remark right after it in [CCGGIIKLLN3]. By our assumption of uniformly bounded curvature and its up to 4-th order derivatives, it is easy to check that, there exists $C_2 > 0$ depending on t_1 and t_2 such that

$$\begin{aligned} &|\nabla L(x)| + |\nabla^2 L(x)| + |\nabla^3 L(x)| + |\nabla^4 L(x)| \le C_2, \qquad x \in \mathbf{M}, \\ &C_2^{-1} L(x) \le d(x, 0, g(t)) \le C_2 L(x), \qquad x \in \mathbf{M}. \end{aligned} \tag{4.19}$$

Here the covariant derivatives are with respect to $g(t)$, $t \in [t_1, t_2]$.

Now, for each $k \ge 0$, let $\lambda_k = \lambda_k(l)$ be a smooth, compactly supported function on $[0, \infty)$ such that $\lambda_k(l) = 1, \quad l \in [0, k]$; $0 \le \lambda_k(l) \le 1, \quad l \in [k, k+1]$; and $\lambda_k(l) = 0, \quad l \in [k+1, \infty)$. We also require $|\lambda_k'| \le 4$. Finally, we take $\phi_k = \lambda_k(L(x))$ as a test function.

By (4.17), we have, since ϕ_k is compactly supported,

$$\begin{aligned} &\frac{d}{dt} \int_{\mathbf{M}} P(u) \phi_k(x) dg(t) = \int_{\mathbf{M}} [\partial_t P(u) - R P(u)] \phi_k(x) dg(t) \\ &= \int_{\mathbf{M}} [\partial_t P(u) - R P(u) + \Delta P(u)] \phi_k(x) dg(t) - \int_{\mathbf{M}} P(u) \Delta \phi_k(x) dg(t) \\ &= \int_{\mathbf{M}} 2\tau \left| Ric - Hess \ln u - \frac{g}{2\tau} \right|^2 u \phi_k(x) dg(t) - \int_{\mathbf{M}} P(u) \Delta \phi_k(x) dg(t). \end{aligned}$$

Let $t_3, t_4 \in [t_1, t_2]$. Integration on the above yields

$$\begin{aligned} &\int_{t_3}^{t_4} \int_{\mathbf{M}} 2\tau \left| Ric - Hess \ln u - \frac{g}{2\tau} \right|^2 u \phi_k(x) dg(t) dt \\ &= \int_{\mathbf{M}} P(u) \phi_k(x) dg(t_4) - \int_{\mathbf{M}} P(u) \phi_k(x) dg(t_3) + \int_{t_3}^{t_4} \int_{\mathbf{M}} P(u) \Delta \phi_k(x) dg(t) dt. \end{aligned}$$

By (4.18), this becomes

$$\begin{aligned} &\int_{t_3}^{t_4} \int_{\mathbf{M}} 2\tau \left| Ric - Hess \ln u - \frac{g}{2\tau} \right|^2 u \phi_k(x) dg(t) dt \\ &\qquad = \int_{\mathbf{M}} i(u) \phi_k(x) dg(t_4) - \int_{\mathbf{M}} i(u) \phi_k(x) dg(t_3) \\ &\qquad\quad - 2\tau \int_{\mathbf{M}} \Delta u \phi_k(x) dg(t_4) + 2\tau \int_{\mathbf{M}} \Delta u \phi_k(x) dg(t_3) \\ &\qquad\quad + \int_{t_3}^{t_4} \int_{\mathbf{M}} i(u) \Delta \phi_k(x) dg(t) dt - 2 \int_{t_3}^{t_4} \int_{\mathbf{M}} \tau \Delta u \Delta \phi_k(x) dg(t) dt. \end{aligned}$$

After integration by parts, we arrive at

$$\begin{aligned} &\int_{t_3}^{t_4} \int_{\mathbf{M}} 2\tau \left| Ric - Hess \ln u - \frac{g}{2\tau} \right|^2 u \phi_k(x) dg(t) dt \\ &= \int_{\mathbf{M}} i(u) \phi_k(x) dg(t_4) - \int_{\mathbf{M}} i(u) \phi_k(x) dg(t_3) + \int_{t_3}^{t_4} \int_{\mathbf{M}} i(u) \Delta \phi_k(x) dg(t) dt \\ &\qquad - 2\tau \int_{\mathbf{M}} u \Delta \phi_k(x) dg(t_4) + 2\tau \int_{\mathbf{M}} u \Delta \phi_k(x) dg(t_3) - 2 \int_{t_3}^{t_4} \int_{\mathbf{M}} \tau u \Delta \Delta \phi_k(x) dg(t) dt. \end{aligned}$$

Notice that the support of $\Delta \phi_k$ is in the region $\{k \le L(x) \le k+1\}$. Since $L(x)$ is comparable with the distance function $d(x, 0, g(t))$, the classical volume comparison theorem tells us that $|\{k \le L(x) \le k+1\}|_{g(t)} \le C e^{ck}$. Now, recall from (4.5) and (4.11) that u

and $i(u)$ have quadratic exponential decay property. Also (4.19) implies that $|\Delta \phi_k| \le C$ and $|\Delta\Delta\phi_k| \le C$. So we can take $\lim_{k\to\infty}$ inside the integrals on the right hand side the last identity. On the other hand, ϕ_k is a nondecreasing function of k, which converges to 1 pointwise. Therefore we can apply the monotone convergence theorem on the left hand side. Therefore

$$\int_{t_3}^{t_4}\int_{\mathbf{M}} 2\tau \left| Ric - Hess \ln u - \frac{g}{2\tau}\right|^2 u dg(t) dt = \int_{\mathbf{M}} i(u) dg(t_4) - \int_{\mathbf{M}} i(u) dg(t_3).$$

This proves the Corollary. □

Finally, we partially extend Perelman's shrinking breather theorem to the noncompact case.

Definition 4.4. *(Breathers)*

A Ricci flow $(\mathbf{M}, g(t))$ *is a called a breather if for some* $t_1 < t_2$ *and* $c > 0$ *there is the relation* $c\phi^* g(t_1) = g(t_2)$ *for a diffeomorphism* ϕ*. The flow in cases* $c = 1$*,* $c < 1$*,* $c > 1$ *are called steady, shrinking and expanding breathers respectively.*

When $\mathbf{M}$ is compact, Perelman [P] proved that a breather is a gradient Ricci soliton, i.e. the Ricci curvature is given by the Hessian of a scalar function. For the noncompact case, we have

Proposition 4.1. *Let* $(\mathbf{M}, g(t))$ *be a noncompact Ricci flow with bounded geometry in the time interval* $[0, T]$*. Suppose* $(\mathbf{M}, g(t))$ *is a shrinking breather in the sense that* $c\,\phi^* g(t_1) = g(t_2)$ *for some diffeomorphism* ϕ*,* $c < 1$ *and* $t_1 < t_2$ *where* $t_1, t_2 \in (0, T)$*. Suppose also* $\mu(g(t_2), \frac{c(t_2-t_1)}{1-c}) < \mu_\infty(g(t_2), \frac{c(t_2-t_1)}{1-c})$*. Then* $(\mathbf{M}, g(t))$ *is a gradient shrinking soliton on the time interval* $[t_1, T]$*.*

Proof.

We follow the same strategy as Perelman's proof for the compact case. The new input is the existence of extremal for the μ invariant in the noncompact setting. Since $\mathbf{M}$ has bounded geometry, we have shown in the proof of Theorem 1.1 (a) that the Log Sobolev functional is bounded from below by a negative constant. By (4.2), the W entropy also has a lower bound for any finite parameter τ. Thus $\mu(g, \tau)$ is a finite number.

Define $L = \frac{t_2 - ct_1}{1-c}$ where c is the number given in the statement of the proposition. Then $c(L - t_1) = L - t_2$. By the scaling and diffeomorphism invariance of the μ invariant, we have

$$\mu(g(t_2), L - t_2) = \mu(g(t_2), c(L - t_1)) = \mu(cg(t_1), c(L - t_1)) = \mu(g(t_1), L - t_1).$$

Note that $L - t_2 = \frac{c(t_2-t_1)}{1-c}$. By the condition $\mu(g(t_2), \frac{c(t_2-t_1)}{1-c}) < \mu_\infty(g(t_2), \frac{c(t_2-t_1)}{1-c})$, we can apply Theorem 1.1 to conclude that $\mu(g(t_2), L - t_2)$ is reached by an extremal function v_2.

Let u be the solution of the final value problem of the conjugate heat equation:

$$\begin{cases} \Delta u - Ru + u_t = 0, \quad t \in [t_1, t_2] \\ u(x, t_2) = v_2^2 \\ \partial_t g(t) = -2Ric, \quad t \in [t_1, t_2]. \end{cases}$$

Since $[t_1, t_2] \subset (0, T)$, Shi's derivative estimate [Sh] shows that the 4-th order derivatives of the curvature tensor are uniformly bounded in $\mathbf{M} \times [t_1, t_2]$. This allows us to use the Corollary. Let $v = v(x,t) = \sqrt{u(x,t)}$. Since v_2 is an extremal of the W entropy at t_2, we know from the Corollary that

$$\begin{aligned}
&\mu(g(t_2), L-t_2) = W(g(t_2), v(\cdot, t_2), L-t_2) \\
&= W(g(t_1), v(\cdot, t_1), L-t_1) + \int_{t_1}^{t_2} \int_{\mathbf{M}} 2\tau \left| Ric - Hess \ln u - \frac{1}{2\tau} g \right|^2 \, u \, dg(t) dt.
\end{aligned}$$

Using $W(g(t_1), v(\cdot, t_1), L-t_1) \geq \mu(g(t_1), L-t_1) = \mu(g(t_2), L-t_2)$, we see that

$$\int_{t_1}^{t_2} \tau \int_{\mathbf{M}} \left| Ric - Hess \ln u - \frac{1}{2\tau} g \right|^2 \, u \, dg(t) dt \leq 0$$

which implies that $Ric - Hess \ln u - \frac{1}{2\tau} g = 0$. i.e. the Ricci flow is a gradient shrinking soliton in the time interval $[t_1, t_2]$. By the uniqueness theorem of Chen and Zhu [CZ] in the noncompact case, the Ricci flow is a gradient shrinking soliton on $[t_1, T]$. This proves the Proposition. □

Acknowledgement. I wish to thank Professor Zhiqin Lu for a useful conversation and Professor Bennett Chow and Professor Lei Ni for their continuous help on my studying of Ricci flow over the years.

References

[Au] Aubin, Thierry, *Problèmes isopérimétriques et espaces de Sobolev.* (French) J. Differential Geometry 11 (1976), no. 4, 573–598.

[BE] Bakry, D.; Émery, Michel, *Diffusions hypercontractives. (French) [Hypercontractive diffusions]* Sminaire de probabilits, XIX, 1983/84, 177-206, Lecture Notes in Math., 1123, Springer, Berlin, 1985.

[BL] Bakry, D.; Ledoux, M. *Lévy-Gromov's isoperimetric inequality for an infinite-dimensional diffusion generator.* Invent. Math. 123 (1996), no. 2, 259-281.

[CCGGIIKLLN2] Chow, Bennett; Chu, Sun-Chin; Glickenstein, David; Guenther, Christine; Isenberg, James; Ivey, Tom; Knopf, Dan; Lu, Peng; Luo, Feng; Ni, Lei, *The Ricci flow: techniques and applications. Part II. Analytic aspects.* Mathematical Surveys and Monographs, 144. American Mathematical Society, Providence, RI, 2008. xxvi+458 pp.

[CCGGIIKLLN3] Chow, Bennett; Chu, Sun-Chin; Glickenstein, David; Guenther, Christine; Isenberg, James; Ivey, Tom; Knopf, Dan; Lu, Peng; Luo, Feng; Ni, Lei, *The Ricci flow: techniques and applications. Part III.* Geometric-analytic aspects. Mathematical Surveys and Monographs, 163. American Mathematical Society, Providence, RI, 2010.

[CGT] Cheeger, Jeff; Gromov, Mikhail; Taylor, Michael, *Finite propagation speed, kernel estimates for functions of the Laplace operator, and the geometry of complete Riemannian manifolds.* J. Differential Geom. 17 (1982), no. 1, 15-53.

[CLY] Cheng, Siu Yuen; Li, Peter; Yau, Shing Tung, *On the upper estimate of the heat kernel of a complete Riemannian manifold.* Amer. J. Math. 103 (1981), no. 5, 1021–1063.

[CN] Jose Carrillo, Lei Ni, *Sharp logarithmic Sobolev inequalities on gradient solitons and applications*, Comm. Analysis and Geometry, 2010.

[CTY] Albert Chau, Luen-Fai Tam and Chengjie Yu. *Pseudolocality for the Ricci flow and applications.* arXiv: math/0701153

[CZ] Chen, Bing-Long; Zhu, Xi-Ping, *Uniqueness of the Ricci flow on complete noncompact manifolds.* J. Differential Geom. 74 (2006), no. 1, 119-154.

[DS] Diaconis, P.; Saloff-Coste, L. *Logarithmic Sobolev inequalities for finite Markov chains*, Ann. Applied Prob. Vol. 6, No. 3, 1996, 695-750.

[F] Federbush, P. *Partially Alternate Derivation of a Result of Nelson* J. Math. Physics, Vol. 10, no 1 Jan. 1969, 50-53.

[G] Gross, Leonard , *Logarithmic Sobolev inequalities.* Amer. J. Math. 97 (1975), no. 4, 1061-1083.

[G2] Gross, Leonard, *Logarithmic Sobolev inequalities and contractivity properties of semigroups.* Dirichlet forms (Varenna, 1992), 54-88, Lecture Notes in Math., 1563, Springer, Berlin, 1993.

[HL] Han, Q. and Lin, F. H. *Elliptic Differential Equations*, Courant Lecture Notes, 1997.

[Heb] Hebey, Emmanuel, *Optimal Sobolev inequalities on complete Riemannian manifolds with Ricci curvature bounded below and positive injectivity radius.* Amer. J. Math. 118 (1996), no. 2, 291–300.

[HV] Hebey, Emmanuel and Vaugon, Michel, *Meilleures constantes dans le thorme d'inclusion de Sobolev.* (French) Ann. Inst. H. Poincar Anal. Non Linaire 13 (1996), no. 1, 57–93.

[L] Lions, P.-L. *The concentration-compactness principle in the calculus of variations.* The limit case. I. Rev. Mat. Iberoamericana 1 (1985), no. 1, 145-201.

[LY] Li, P.; Yau, S.T., On the parabolic kernel of the Schrödinger operator, *Acta Math.*, 156 (1986) 153-201.

[OV] Otto, F.; Villani, C. *Generalization of an inequality by Talagrand and links with the logarithmic Sobolev inequality.* J. Funct. Anal. 173 (2000), no. 2, 361-400.

[P] Perelman, Grisha, *The entropy formula for the Ricci flow and its geometric applications*, Math. ArXiv, math.DG/0211159.

[Rot] Rothaus, O. S. *Logarithmic Sobolev inequalities and the spectrum of Schrdinger operators.* J. Funct. Anal. 42 (1981), no. 1, 110-120.

[Sa] Saloff-Coste, Laurent, *Uniformly elliptic operators on Riemannian manifolds.* J. Differential Geom. 36 (1992), no. 2, 417–450.

[Sh] Shi, Wan-Xiong, *Deforming the metric on complete Riemannian manifolds.* J. Differential Geom. 30 (1989), no. 1, 223-301.

[Z] Zhang, Qi S., *Sobolev inequalities, heat kernels under Ricci flow and the Poincaré conjecture*, CRC Press 2011.

e-mail: qizhang@math.ucr.edu

Department of Mathematics, University of California, Riverside, CA 92521, USA